Resistance and **Tolerance to Citrus Greening Disease AKA Huanglongbing** or HLB Ed Stover- USDA/ARS Ft. Pierce, FL



Support Scientists and Postdocs Randy Driggers Guixia Hao Sharon Inch Malu Oliveira Ric Stange Godfrey Miles



Principle Collaborators: **David Hall Greg McCollum Bob Shatters** YongPing Duan Goutam Gupta **Gloria Moore Richard Lee** Manjunath Keramane Chandrika Ramadugu **Bill Belknap Jim Thomson Fred Gmitter** Jude Grosser **Mikeal Roose** Malcolm Smith

Huanglongbing, AKA Citrus Greening

- Caused by a bacterium, Liberibacter asiaticus (Las), which in trees lives only in the plumbing which distributes sugar throughout the plant (called the phloem)
- Transmitted in the US only by an insect that feeds on citrus leaves (and specifically the phloem) called the Asian Citrus Psyllid (ACP), much as mosquitoes carry the malaria pathogen
- Most distinctive symptom is blotchy mottle of the leaves in which the yellow areas are not symmetrical on either side of the leaf midrib (see below). In severely infected trees, many fruit are lopsided with aborted seeds and poor color development (also below)
- Within a few years of infection, many citrus trees become weak, have poor quality fruit, with lots of fruit drop, and trees may die or become useless

Photos Bové, 2006





HLB tolerance/resistance solutions in short-, medium-, and long-term

- Existing cultivars and most advanced selections
- New selections with conventional citrus cultivar genetics
- New hybrids entering testing
- Hybrids with more-resistant citrus relatives
- Transgenics- best chance of total immunity

All these solutions require replanting! We need therapies to maintain production in existing trees to provide cash flow for new plantings.

Categories of resistance:


Immune- is the best! No disease whatsoever. Example: humans (and most plants) are immune to HLB! No good evidence of HLB-immune citrus

<u>Resistant</u>- slower to develop infection and/or slower to develop symptoms and/or lower levels of the pathogen result; also lower levels of symptoms

<u>Tolerant</u>- some symptoms and even the pathogen may develop at levels similar to those in susceptible individuals/genotypes. However, they continue to grow (and produce) fairly normally.

>Example: sweet orange is tolerant to sour orangedecline CTV on appropriate stock

- In time truly immune trees will be found... in the meantime
- How much of a delay in symptom development / compromise of cropping is needed to be useful?




Overview of US Citrus

- In US: total value of citrus industry is ~\$9 billion -75% production is sweet orange -11% grapefruit
- FL: 66% of US Citrus total -85% oranges (96% juiced) -12% grapefruit (58% juiced)

Citrus #1 fruit consumed in US

50x10⁶/ yr cartons fresh exports



Florida Citrus Statistics 2011-2012 (February 2013) USDA, National Agricultural Statistics Service

Fruit by % US Consumption

FL and US Citrus is almost a monoculture

•All sweet oranges are mutants of a hybrid that arose in China a thousand or more years ago, and so are almost genetically identical

•Commercial grapefruit are similarly mutants of a hybrid (between sweet orange and pummelo) that arose in the Caribbean a few hundred years ago

•The genetic homogeneity of US citrus provides great vulnerability to introduced pathogens or pests

•As it happens, sweet orange and grapefruit display among the most severe decline from HLB of any citrus tested!


Slide 8

gm1 gmccollum, 5/28/2014

gm2 gmccollum, 5/28/2014

Significant tolerance to HLB in existing cultivars? Survey in groves with multiple types in 2010 -disease introduced into trees at maturity Liberibacter per sample by cultivar Stover & McCollum

	Pathogen titer: mean # CLas	% trees
	/100 mg sample by PCR	"HLB+"
Minneola	304	43%
Murcott	168	44%
Sweet orange	236	31%
Grapefruit	40	20%
Temple	9	15%
Fallglo	13	18%
Sunburst	107	13%



Grapefruit vs. Near Grapefruit

	3 yr C	umu		201				
	Fruit		Fruit		Disease		TSS/TA	
Cultivar	per tree		drop		rating		ratio	
Flame	129.4	bc	50%	b	4.2	b	7.0	b
Marsh	66.5	С	53%	b	4.4	b	5.7	С
Jackson	219.9	ab	14%	а	2.5	а	10.6	а
Triumph	255.1	а	15%	а	2.4	а	9.6	а
F&M vs. T&J	0.0002		<0.0001		< 0.0001		0.0001	



- Fruit quality assessments were made each growing season with 'Triumph'/'Jackson' showing generally acceptable commercial fruit quality
- 'Flame'/'Marsh' had too low Brix/acid.
- In 2011/2012 many 'Flame'/'Marsh' were small and/or misshapen while 'Triumph'/ 'Jackson' displayed normal size and shape.
 More evidence of
- Similar levels of HLB bacterium tolerance

Ongoing Evidence that some Mandarins have substantial HLB



Kunembo (C. nobilis)

Dancy

What if trees are exposed to CLas at planting? > 6 yr replicated trial at Picos Farm >CLas titers not significantly different

			Fruit/tree		Health		Change	in
Scion/Rootstock	Mortality (%)		<u>Oct 2015 (no.)</u>		<u>Oct 2015 (3 pt)</u>		<u>diam. (mm)</u>	
Fallglo/Kinkoji	20	а	28.4	b	1.9	cd	23.8	b
Hamlin/Cleopatra	20	а	18.6	bc	2.2	bc	20.4	b-d
Hamlin/Kinkoji	10	а	12.9	cd	1.9	cd	14.5	d
Ruby/Kinkoji	10	а	4.6	е	1.6	d	20.7	bc
SugarBelle/Sour	0	а	81.3	а	2.9	а	46.1	а
Tango/Kuharske	0	а	88.1	a	2.9	а	32.2	а
Temple/Cleopatra	18	а	35.6	a 🗸	2.3	ab	23.8	b

 Some scion/rootstock combinations continued to develop even with high titers of CLas and and strong mottle symptoms
Not "tolerant" rootstocks used so likely a scion effect



SugarBelle/SourOr and Tango/Kuharske look particularly good and are producing more fruit

Hamlin/Kinkoji

SugarBelle/Sour Orange

Tango/Kuharske







5-51-2 Clementine x Orlando cross by J. Hearn

Several USDA C x O are displaying potentially useful HLB tolerance

New planting looking at wide range of Clem x Orlando to map tolerance genes- Stover & Roose



We are most importantly a breeding program, and make >2000 new hybrids each year. All new hybrids from crosses intended to achieve HLB tolerance/ resistance are planted at the Picos farm in Ft. Pierce



It's important to know how our new selections hold up to HLB! Replicated plants of 50 USDA selections and standards planted in the field after no-choice hot-ACP for a week and 4 months in the hot-ACP house



New "Sweet Orange" Hybrids

-different genetic makeup than all other sweet oranges- MAY have greater resistance/tolerance and currently being tested -Have been propagated and planted in commercial field tests (McCollum lead)

Finally making progress on truly sweet-orange-like hybrids!



FF-1-75-55 Ambersweet x FF-1-30-52. Ripens about the same time as Hamlin, has orange appearance, taste, and aroma volatiles. Can be peeled by hand. Near Valencia Juice Color. Relatively few seeds. Easy peeling

Volatile profile comparison of USDA sweet-orange-like hybrids vs. 'Hamlin' and 'Ambersweet'



Jinhe Bai, Elizabeth Baldwin Randall Driggers, Jack Hearn and Ed Stover

Further afield-Considerable HLB resistance in citrus gene pool! Ramadugu et al. 2016

 Field experiment with genebank at Riverside CA of 85 citrus relative genotypes - showed *Poncirus* among most-resistant to HLB and also psyllid colonization (ARS CA and FL)

 Eremocitrus and Microcitrus, also showed strong Las and psyllid resistance and we have new collaboration with Queensland citrus breeder Malcolm Smith









Trifoliate genes for HLB resistance

•U of Florida (Fred Gmitter), UC Riverside (Roose) and USHRL (Stover) collaborating on trial to identify genes associated with HLB-resistance in citranges

•When mapped and identified, can use gene markers in conventional breeding and in intragenics

 Includes near commercial quality, advanced Poncirus hybrids



U1 REd text: Confusing USHRL, 6/2/2010

Poncirus is deciduous, cold-hardy, Citrus crosscompatible, but tastes terrible

USDA started using *Poncirus* as parent 110 yrs ago for coldhardiness: looks like may pay off for HLBtolerance


Sweet Orange like fruit-Navel in alligator-hide Apparent tolerance to HLB 1/16 *Poncirus* No off-flavor Being used in many crosses 100+ Poncirus hybrid genotypes all replicated and exposed to HLB/ACP for 36 months

Gnarlyglo trees are the largest (7 ft vs 3 ft in sweet orange) healthiest trees in the entire planting, even though produced from field budwood



Transgenics for HLB- Resistant Citrus

- Tolerance and resistance is great..... IF you have decided to live with HLB
- Transgenics appear to be the most promising solution for strong HLB resistance and perhaps <u>immunity</u>
- Another major advantage is ability to improve an existing cultivar with essentially no other changes: HLB-resistant Valencia, Hamlin, Ray Ruby etc.



Future of transgenic Citrus

•Genetically engineered (GE) cultivars deregulated for commercial use in ~25 different agricultural crops

•GE crops are grown on ~12% of global arable land, mainly four field crops: soybean, maize, cotton and canola

•Several GE horticultural crops are being produced commercially since they provide solutions to otherwise intractable threats, much as HLB seems for citrus


 Commercial GE citrus is likely inevitable and GE crop concerns will likely decline with time

NO released transgenic in any crop for bacterial resistance

Categories: Transgenic Strategies


- Direct attack on the pathogen
 - >Antimicrobials
 - >"Antibodies" for exposed pathogen proteins
- Host pathogen interactions
 - >Basal defense genes
 - >CLas gene products that target host (nuclear localization protein, flagellin etc.)
- Citrus physiology
 - >Possibly overactive defense response
- Psyllid targets








Thionin and chimeral antimicrobial peptides, designed by Goutam Gupta (Los Alamos National Laboratory)



Chimera of a citrus serine protease (cyan) joined to the lytic D4E1 peptide (red) by a GSTA linker (yellow)

Xcc Infiltration results with transgenic plants containing thionin, D4E1 and chimera




Transgenic Carrizo grafted with HLB+Rough lemon- in progress






Aggressive challenge begins with no-choice exposure to CLas infected psyllids- led by D. Hall



Trees in greenhouse with free-flying CLas infected psyllids plus source plants-led by D. Hall

Spinach Defensins for HLB resistance: Erik Mirkov Texas A&M University and Southern Gardens Citrus



•Furthest along in deregulation

•EUP (experimental use permit) is in place for transgenic trees expressing two spinach defensins


- Can plant up to 400 ac in Florida and 200 ac in Texas

•Temporary tolerance exemption has been granted by the EPA for two spinach defensins in GMO trees – but will not use in commerce yet

•EUP has been submitted for the expression of spinach defensins using the CTV viral vector as the delivery method

- Does not result in a GMO tree; Can plant up to 400 ac in Florida

•Temporary tolerance exemption has been submitted for the expression of spinach defensins using the CTV viral vector- but won't use in commerce yet





A Race to Save the Orange by Altering Its DNA



Trees that are infected by disease are cut down and burned in Clewiston, Fla., at groves owned by Southern Gardens Citrus. Richard Perry/The New York Times

🔛 E-MAIL	By AMY HARMON JULY 27, 2013 ₹ 776 COMMENTS
FACEBOOK	CLEWISTON, Fla. – The call Ricke Kress and every other citrus grower in Florida
	dreaded came while he was driving.
PRINT	"It's here" was all his grove manager needed to say to force him over to the side of the



Carl Külsgaard/ZUMA

In a July 27 feature article that set the interwebs aflame, New York Times reporter Amy Harmon told the tale of a bacterial pathogen that's stalking the globe's citrus trees, and a Florida orange juice company's effort to find a solution to the problem through genetic engineering.

An invasive insect called the Asian citrus psyllids carries the bacteria, known as Candidatus Liberibacter, from tree to tree, and it causes oranges and other citrus fruits to turn green and rot. "Citrus greening," as the condition has become known, has emerged as a pest nearly wherever citrus is grown globally. Harmon reported that an "emerging scientific consensus" holds that only genetic engineering can defeat it.

Meanwhile, Michael Pollan, a prominent food industry and agribusiness critic, tweeted this:

The FUTURE for Citrus production looks bright!

- Still many dark days to get there, at least in Florida
- Massive research investment is generating options for future plantings with HLB-resistant/tolerant trees

Research is poised to identify game-changing technologies to protect uninfected susceptible trees such as ability to quickly identify and remove newly infected trees or prevent ACP from carrying CLas

•Acceptance of "replacement" citrus varieties or transgenics may be critical for an HLB solution

•An important missing piece in FL is therapy to maintain production on existing trees, needed to provide cash for implementation of new solutions

Thanks!

- Florida Citrus Research & Development Foundation
- <u>NIFA- Specialty Crops Research Initiative</u>
- New Varieties Development and Management Corp
- Florida Citrus Research Foundation (Whitmore)
- California Citrus Research Board
- DPI Budwood Office (especially Peggy Sieburth)
- USDA/ARS Funding and USDA/APHIS

Jodi AvilaRobyn BaberWayne BrownScott CilientoJacqueline DepazBelkis DiegoAmber HollandDiane HelsethChris LasserDavid LindseySteve MayoKathy MoultonLuc OverholtSean ReifJames SalvatoreMatthew SewellRegina TracyAshley Witkowski

Abby Bartlett Ellen Cochrane Lynn Faulkner Scott Hyndman Philip Matonti Jerry Mozoruk Mike Rutherford Jeff Smith Jen Wildonger

