

Hendry County Extension, P.O. Box 68, LaBelle, FL 33975 (863) 674 4092

Flatwoods Citrus



Vol. 20, No. 5

May 2017

Dr. Mongi Zekri
Multi-County Citrus Agent, SW Florida



Table of Contents

Upcoming Events	2-5
Flatwoods Citrus Newsletter Sponsors – Thank you!	6-9
El Niño/Southern Oscillation (ENSO) Diagnostic Discussion	10
Implementing BMPs	11
Neutralizing Excess Bicarbonates from Irrigation Water in Florida	12-15
Frequently Asked Questions About Biosolids	16-20
Citrus Rust Mites	21-23
Greasy Spot Fungal Disease	24-25

The Twenty-Seventh Annual Farm Safety Day

Friday, 12 May 2017
Saturday, 13 May 2017

AN IMPORTANT MESSAGE TO EMPLOYERS

Safe and competent equipment operators are important to you as an employer. Accidents, which cause damage, injury or death to employees, equipment and crops, are costly. We believe all types of accidents can be reduced with proper employee training. Our training has been designed to help your employees perform better, operate safely to prevent accidents, fulfill necessary training requirements and build pride in themselves and their farm company.

Certificates

The 2017 Southwest Florida Farm Safety Day is almost here. Farm Safety Day is an educational event designed to emphasize the importance of farm/equipment safety. Each participant is presented with a certificate of attendance and **the employer will be provided with a certificate of training that can be placed into the employee's file.**

Sessions this year:

1. WPS for Handlers
2. Safety Culture
3. Respiratory Protection
4. Health Hazards

Certified pile burners class July 12, 2017

Pre-registration is required to attend, and class size is limited to the first 50 people.

Registration fee: \$50

The \$50 fee covers the training sessions, a booklet with all the presentations in color, other handouts, refreshments, and lunch.

PRE-REGISTRATION WILL NOT BE ACCEPTED WITHOUT PAYMENT OF THE REGISTRATION FEE.

Send your registration form and check as soon as possible. **Only 6 seats are available. Call (239 595 5494) and reserve your seat before sending your registration form and check.**

Location: The Immokalee IFAS Center

The Florida Division of Forestry and University of Florida Cooperative Extension Service will be conducting a Certified Pile Burners Course that will show you how to burn piles **legally, safely and efficiently.**

Most importantly, it could save a life. This training will be held from 8:00 am till 4:30 pm at the **Southwest Florida Research and Education Center in Immokalee.**

Detailed information is available online:

http://www.freshfromflorida.com/content/download/74623/2126893/Certified_Pile_Burner_Hendry_July12-2017.pdf

All You Need to Know About Scouting and Managing Citrus Canker and Citrus Fungal Diseases

Date & time: Wednesday, May 10, 2017, 10:00 AM – 12:15 PM

Location: UF-IFAS Southwest Florida Research and Education Center, Immokalee

Speakers: Dr. Megan Dewdney and Dr. Evan Johnson, UF-IFAS

Program Coordinator: Dr. Mongi Zekri, UF-IFAS

Program Sponsor: Doug Wilbanks with Syngenta

2 CEUs for pesticide license renewal

2 CEUs for certified crop advisors (CCAs)

Pre-registration is required. No registration fee and lunch is free Thanks to **Doug Wilbanks with Syngenta.** To reserve a seat, call 863 674 4092, or send an e-mail to Dr. Mongi Zekri at: maz@ufl.edu

Agenda

----10:00 AM – 10:55 AM

1. Scouting and managing melanose, greasy spot and citrus black spot (Causal agent, disease cycle, cultural control, leaf litter, chemical control)

Dr. Megan Dewdney, UF-IFAS

10:55 AM – 11:05 AM Break

----11:05 AM - 12:00 Noon

2. Scouting and managing citrus canker and Phytophthora root rot, foot rot, and brown rot

(cultural practices, preventive control, sampling, timing for chemical control, chemical products)

Dr. Evan Johnson, UF-IFAS

----12:00 Noon – 12:15 PM

3. Syngenta Citrus Product Update
The New Insecticide (Minecto Pro)

Doug Wilbanks, Syngenta



Citrus blight, leprosis, Diaprepes and nematode management

Date & time: Wednesday, May 24, 2017, 10:00 AM – 12:00 Noon

Location: UF-IFAS Southwest Florida Research and Education Center, Immokalee

Speakers: Dr. Larry Duncan and Dr. Ron Brlansky, UF-IFAS

Program Coordinator: Dr. Mongi Zekri, UF-IFAS

Program Sponsor: Joe Mitchell with BASF

2 CEUs for pesticide license renewal

2 CEUs for certified crop advisors (CCAs)

Pre-registration is required. No registration fee and lunch is free Thanks to **Joe Mitchell with BASF**. To reserve a seat, call 863 674 4092, or send an e-mail to Dr. Mongi Zekri at: maz@ufl.edu

Agenda

----10:00 AM – 10:55 AM

1. Diaprepes and nematodes

(Root weevils, nematodes, background, fumigation, cultural practices, sanitation, cultural management, chemical management)

Dr. Larry Duncan, UF-IFAS


10:55 AM – 11:05 AM Break

----11:05 AM - 12:00 Noon

2. Citrus blight and leprosis

(transmission, symptoms, distribution, host range, causal agent, diagnosis)

Dr. Ron Brlansky, UF-IFAS



Citrus Undercover Production Systems (CUPS) and Controlled-Release Fertilizers (CRFs)

Date & time: **Wednesday, June 28, 2017, 10:00 AM – 12:00 Noon**

Location: UF-IFAS Southwest Florida Research and Education Center, Immokalee

Speakers: **Dr. Arnold Schumann, Pete Spyke, Ward Gunter, and Dr. Nael El-Hout**

Program Coordinator: Dr. Mongi Zekri, UF-IFAS

Program Sponsor: **Ward Gunter, ICL Specialty Fertilizers**

2 CEUs for pesticide license renewal

2 CEUs for certified crop advisors (CCAs)

Pre-registration is required. No registration fee and lunch is free Thanks to **Ward Gunter, ICL Specialty Fertilizers.** To reserve a seat, call 863 674 4092, or send an e-mail to Dr. Mongi Zekri at: maz@ufl.edu

Agenda

----10:00 AM – 10:50 AM

1. Citrus undercover production systems (CUPS)

Dr. Arnold Schumann, UF-IFAS

10:50 AM – 11:00 AM Break

----11:00 AM - 11:50 AM

2. Controlled-release fertilizers (CRFs) for citrus

Pete Spyke, Arapaho Citrus Management, Inc.

----11:50 AM - 12:20 PM

3. Maximizing the efficiency of citrus nutritional programs

Ward Gunter, Sales Lead & Business Development Manager, ICL Specialty Fertilizers



Special Thanks to sponsors of the "Flatwoods Citrus" newsletter for their generous contribution and support. If you would like to be among them, please contact me at 863 674 4092 or maz@ufl.edu



Sam Thayer
President

P.O. Box 1849
Dundee, FL 33838
Phone: (863) 439-3667
Fax: (863) 439-6608

www.maxijet.com
sam@maxijet.com

The Standard of Quality In Low-Volume Irrigation

Donald Allen
AGLIME SALES, INC.
1375 Thornburg Road
Babson Park, FL 33827-9549
Mobile: 863 287 2925
Agnnet # 52925
donald.allen@aglimesales.com

Steve Fletcher
Fletcher Flying Service, Inc.
Phone: 239 860 2028
Fax: 863 675 3725

Scott Houk
Dow AgroSciences
13543 Troia Drive
Estero, FL 33928
Phone: 239-243-6927
SEHouk@dow.com



G.P. SOLUTIONS™

We've Got You Covered From The Bottom To The Top

- Root Health Programs
- Foliar Nutrient Programs
- Disease Control Programs

Liquid Fertilizers, Micronutrients & Natural Organics

900 Cowboy Circle, Labelle, FL 33935
(863) 675-1500 | www.Nitro30.com

Ed Early
DuPont Crop Protection
P O Box 7768
Fort Myers, FL 33911
Phone: 239-994-8594
Edward.L.early@dupont.com

Jim Faircloth
syngenta®
Phone: 402 639 2756
jim.faircloth@syngenta.com



**BLACKSMITH
BIOSCIENCE**

Jack Kilgore
Technical Sales Rep, SE US

Office: 7150 East Brentwood Road
Ft. Myers, FL 33919

(239) 707-7677
g8trmanjek@comcast.net
www.blacksmithbio.com

Billy Hopkins
Hopkins Nursery
239 658 0370
tropicals@wildblue.net
Tropical fruit & peach trees




Bo Meador
863 673 0727
www.AgroSource.net
bo.meador@agrosource.net

NICHINO AMERICA
Scott Croxton
scroxton@nichino.net
Samuel S. Monroe
smonroe@nichino.net
www.nichino.net



Nufarm
Craig Noll
Cell: 239 549 4474
Craig.Noll@us.nufarm.com



YaraLiva[®]
A whole lot more than calcium nitrate.

Eric Waldo, Farmer Engagement Manager,
352-215-8480, eric.waldo@yara.com

Richard Newman, Regional Sales Manager,
904-923-9595 richard.newman@yara.com

Arysta LifeScience
Shaun Yule
Phone: 386-561-0493
E-mail: shaun.yule@arysta.com



Magna-Bon[®]
AGRICULTURAL CONTROL SOLUTIONS

Frank Miele
Office: 863 357 0400
Cell: 954 275 1830
Fax: 863 357 1083
E-mail: famiele1@aol.com



Plant Food Systems, Inc.

P.O. Box 775
Zellwood, FL 32798
Tel: 407 889 7755

Special Thanks to sponsors of the "Flatwoods Citrus" newsletter for their generous contribution and support. If you would like to be among them, please contact me at 863 674 4092 or maz@ufl.edu



Imidan
Agricultural Insecticide

Bart Hoopingarner
(941) 737-7444
bhoopingarner@gowanco.com



**PHILLIP RUCKS
CITRUS
NURSERY**

Citrus, Peaches, Windbreaks
Phone: 863 635 1948
E-mail: info@rucksnursery.com
www.ruckscitrusnursery.com

Your Best Source for Florida Land Values!



SREland.com/market-reports
877-518-LAND



Saunders
REAL ESTATE

Heath Prescott

KeyPlex[®]

Toll Free: 800 433 7117
Mobile: 863 781 9096
Nextel: 159*499803*6

Sarah Markle
863-673-8699

Sarah.Markle@valent.com



Wellmark

Extinguish
PLUS

EXTINGUISH
Professional Fire And Heat

www.extinguishfireants.com

Jack Kilgore

M: 239-707-7677

g8trmanjek@comcast.net

Michael Seese

239 336 9152

mpseese@aol.com

Infinity Ag Solutions

infinityagsolutions.com



SPECIALIZING IN LARGE ACREAGE AGRICULTURAL and TRANSITIONAL LAND

Land Solutions inc.

AGRICULTURAL
LAND SERVICES
More Than Just Dirt.
www.landsolutions.net
239-489-4066

Adrian Jahna
BASF Corporation

Cell: 863 443 2404

Adrian.jahna@basf.com

Sim Nifong



SENifong@timacusa.com

Cell: 863 441 1057

<http://www.timacusa.com>

FMC

FMC Corporation APG

Eric Johnson

Lakeland, FL

352-281-2325

EJ.Johnson@FMC.Com

fmccrop.com

actagro.

Reese Martin

Reese.Martin@actagro.com

863 605 8533

www.actagro.com

AgroFlite

"Ultimate Precision in Aerial Application"

CUSTOM AERIAL APPLICATION

Helicopter & Fixed-Wing Aircraft

Call for our Discounted Pricing today

(561) 600-8885

www.agroflite.com

Special Thanks to sponsors of
the "Flatwoods Citrus"
newsletter for their generous
contribution and support. If
you would like to be among
them, please contact me at
863 674 4092 or maz@ufl.edu

EL NIÑO/SOUTHERN OSCILLATION (ENSO) DIAGNOSTIC DISCUSSION

issued by

**CLIMATE PREDICTION CENTER/NCEP/NWS
and the International Research Institute for Climate and Society**

13 April 2017

ENSO Alert System Status: Not Active

Synopsis: ENSO-neutral conditions are favored to continue through at least the Northern Hemisphere spring 2017, with increasing chances for El Niño development by late summer and fall.

ENSO-neutral conditions continued during March, with near-average sea surface temperatures (SSTs) across the central equatorial Pacific and above-average SSTs in the eastern Pacific (Fig. 1). The latest weekly Niño index values were near zero in the Niño-4 and Niño-3.4 regions, and +0.8 and +0.9°C farther east in the Niño-3 and Niño-1+2 regions, respectively (Fig. 2). The upper-ocean heat content anomaly, averaged across the central and eastern Pacific (Fig. 3), decreased to near zero during March, a reflection of above-average temperatures at depth in the east offset by below-average temperatures in the central Pacific (Fig. 4). Atmospheric convection remained suppressed over the central tropical Pacific and enhanced over the Maritime Continent (Fig. 5). The low-level easterly winds were enhanced over the central and western tropical Pacific, and weaker than average over the eastern Pacific. Also, upper-level westerly winds were anomalously easterly over the western and far eastern Pacific, while the Southern Oscillation Index was near average. Overall, the ocean and atmosphere system is consistent with ENSO-neutral conditions.

Most models predict the continuation of ENSO-neutral (3-month average Niño-3.4 index between -0.5°C and 0.5°C) through the late Northern Hemisphere spring (April-June; Fig. 6). However, at least one-half of the dynamical model forecasts, including the NCEP CFSv2, anticipate an onset of El Niño as soon as the April-June season. Because of typically lower skill in forecasts made at this time of the year, and the lingering La Niña-like tropical convection and wind patterns over the western half of the Pacific basin, the forecaster consensus favors ENSO-neutral during April-June with a 60-65% chance. Thereafter, there are increasing odds for El Niño toward the second half of 2017 (~50% chance from approximately August-December). In summary, ENSO-neutral conditions are favored to continue through at least the late Northern Hemisphere spring 2017, with increasing chances for El Niño development by late summer and fall (click [CPC/IRI consensus forecast](#) for the chance of each outcome for each 3-month period).

This discussion is a consolidated effort of the National Oceanic and Atmospheric Administration (NOAA), NOAA's National Weather Service, and their funded institutions. Oceanic and atmospheric conditions are updated weekly on the Climate Prediction Center web site ([El Niño/La Niña Current Conditions and Expert Discussions](#)). Forecasts are also updated monthly in the [Forecast Forum](#) of CPC's Climate Diagnostics Bulletin. Additional perspectives and analysis are also available in an [ENSO blog](#). The next ENSO Diagnostics Discussion is scheduled for 11 May 2017. To receive an e-mail notification when the monthly ENSO Diagnostic Discussions are released, please send an e-mail message to: ncep.list.enso-update@noaa.gov.

Climate Prediction Center
National Centers for Environmental Prediction
NOAA/National Weather Service
College Park, MD 20740

What Are Agricultural Best Management Practices?

Agricultural **Best Management Practices** (BMPs) are practical measures that producers can take to reduce the amount of fertilizers, pesticides, animal waste, and other pollutants entering our water resources. They are designed to improve water quality while maintaining agricultural production. The Florida Department of Agriculture and Consumer Services (FDACS) has adopted BMPs for most commodities in the state. Each BMP manual covers key aspects of water quality and water conservation. Typical practices include:

Nutrient Management to determine nutrient needs and sources, and manage nutrient applications (including manure) to minimize impacts to water resources.

Irrigation Management to address the method and scheduling of irrigation to reduce water and nutrient losses to the environment.

Water Resource Protection using buffers, setbacks, and swales to reduce or prevent the transport of sediments and nutrients from production areas to waterbodies.

Why Should I Implement BMPs?

- Implementing (and maintaining) verified FDACS-adopted BMPs provides a presumption of compliance with state water quality standards for the pollutants addressed by the BMPs.
- Some BMPs can help you operate more efficiently and reduce costs, while you help protect the environment.
- Producers who implement FDACS-adopted BMPs might satisfy some water management district (WMD) permitting requirements. Check with your WMD.
- With some exceptions, the Florida Right to Farm Act prohibits local governments from regulating an agricultural activity that is addressed through rule-adopted BMPs that producers are implementing.
- The Florida Department of Environmental Protection is developing Basin Management Action Plans (BMAPs) to meet adopted water quality targets called Total Maximum Daily Loads (TMDLs). *Where FDEP adopts a BMAP that includes agriculture, producers must either implement FDACS-adopted BMPs, or conduct monitoring (prescribed by FDEP or the water management district) to show they are not violating water quality standards.* This type of monitoring is very expensive.

How Do I Participate in BMPs?

1. **Schedule a meeting with a BMP team member**, who will provide a free FDACS BMP manual and other BMP-related information.
2. **Participate with the coordinator in a free assessment of your operation** to determine which BMPs apply to you.
3. **Fill out a BMP checklist and sign the Notice of Intent (NOI)** to implement the BMPs.
4. **Keep a copy** of the checklist and signed NOI in your records.
5. **Implement and maintain the applicable BMPs and keep adequate records** to maintain a presumption of compliance with state water quality standards.

Contact Information

Dr. Kelly Morgan
Phone: (239) 658-3413
conserv@ufl.edu

If you would like to receive a Certificate of Enrollment in BMPs:

Contact FDACS at: (850) 617-1727 or
AgBmpHelp@FreshFromFlorida.com

Neutralizing Excess Bicarbonates from Irrigation Water in Florida

By Gerald Kidder and Ed Hanlon, UF-IFAS



Many sources of irrigation water in Florida contain dissolved bicarbonates. Irrigation with such water can cause adverse plant growth by excessively raising the pH of the soil. The magnitude of the effect depends on the concentration of the bicarbonates in the water, the amount of the water applied, the buffering capacity of the soil, and the sensitivity of the citrus variety/rootstock being grown.

This publication addresses this important water quality problem and suggests management practices to minimize adverse effects on citrus tree growth and production.

1. Where in Florida is the problem most likely to occur?

The problem of high dissolved bicarbonates is likely to occur wherever water comes from a limestone aquifer, such as the Floridan or Biscayne, or from lakes or canals that cut into limestone. Thus, this is a potential problem in most of Florida.

2. How can I find out if I have high-bicarbonate water?

A water test is the surest means of determining if a problem exists.

3. Isn't it sufficient to just measure the water's pH?

If the pH of your irrigation water is below 7.0, then we may safely assume that it will not be a significant source of liming materials. However, if the pH is above 7.0, we know that the water contains bases but we don't know how much. For example, one water source may have a relatively high pH of 8 and yet contain a very low level of bicarbonates. Another water source, with the same pH, may have a very high bicarbonate level.

4. How are Ca and Mg analyses useful?

Multiplication of parts per million (ppm) Ca by 0.05 and ppm Mg by 0.083, and summing the two products, will give the milliequivalents of those cations per liter (me/L) of water. In many cases, Ca and Mg will be associated with bicarbonate and carbonate salts. Under those conditions, the me/L of Ca plus Mg will be a good estimate of the me/L of associated bases. However, if other non-basic ions such as sulfate are present, the calculation would overestimate the base content of the water. Thus, Ca and Mg analyses may be useful in estimating base content but should be used with caution.

5. In which crop situations am I likely to have a problem with high pH water?

Trifoliolate and most trifoliolate hybrid rootstocks are particularly sensitive to high pH soil, are trees budded onto them usually exhibit ill effects of high bicarbonate water through micronutrient deficiency symptoms. Trees budded on Swingle rootstock are well-known for their sensitivity to pH-induced iron chlorosis. Trees budded on citrange rootstocks have shown manganese and zinc deficiencies when the soil pH has been raised by heavy or prolonged use of "hard" water (i.e., water with lots of Ca and Mg bicarbonates).

6. Which irrigation situations are most problematic?

Heavy irrigations applied to soils of low buffering capacity will present the most problems to citrus trees.

7. What can I do to minimize the adverse effects of high-bicarbonate water?

Be careful not to over-irrigate. Know the water holding capacity of your soil and apply only enough water without exceeding the root zone water-holding capacity. Over-irrigation is costly in many ways -- the cost of pumping, of leached nutrients, of wasted water resources and, in this case, of accelerating the increase in soil pH. Avoid these with good irrigation management.

Apply acids or acid-forming materials to the soil to counteract the bases applied in the water.

Neutralize the liming effect of the water by adding acid to the water before it is applied to the trees.

8. What can be done if the trees are already suffering from water-induced high pH?

Where high levels of bicarbonates in the water have caused soil to be too high for proper tree performance, it may be necessary to lower the soil pH. This may be accomplished by addition of extra acid in the irrigation water, use of acid-forming fertilizer in certain cases, or application of elemental sulfur to the soil.

It is important to note that the acid-producing effect of sulfur comes from the formation of sulfuric acid when soil bacteria act on the elemental sulfur. The sulfate form of sulfur applied in fertilizers such as potassium sulfate, magnesium sulfate, or gypsum (calcium sulfate) does not have the acid-producing effect of elemental sulfur.

Sulfur application rates of 300 to 500 pounds per acre should not be exceeded. This rate is equivalent to between 0.7 and 1.1 lbs/100 square feet of treated surface area. Over-application of sulfur or acid

can cause damage to trees, an effect you certainly want to avoid. Monitor changes carefully.

Remember the pH will increase again as you continue to irrigate with high bicarbonate water. Water or soil acidification will be a continuing effort.

9. Can acid-forming fertilizers keep the soil pH from getting too high?

Under many circumstances, the quantity of bases that is being supplied in the irrigation water far exceeds the quantity of acid formed by addition of fertilizers. Under those conditions acid-forming fertilizer will not control the problem of increasing soil pH.

10. How can I neutralize the bicarbonates in my irrigation water?

Injection of acid into the irrigation water is a direct way of neutralizing the bases present. Acid may be injected in much the same way as fertilizer. You must take precautions to avoid injuring yourself and your trees and to avoid contamination of the aquifer. These points are discussed below.



11. How much acid should I apply?

The amount of acid that you mix with the irrigation water will depend on the quantity of bases your water contains and on the strength of the acid you use. The base content of the water is determined in the water test and the strength of the acid is given on the container. One milliequivalent (me) of acid completely neutralizes one milliequivalent of base. For example, if an irrigation water contains 5.2 me of bases per liter, it would take 5.2 me of acid to completely neutralize the liter of water. Neutralization

of 80 to 90% of the bases in water is a reasonable goal for most irrigation situations.

Multiply the factor by the milliequivalents of base per liter (me/L) which your water contains. This value is determined in the laboratory test of your water or is estimated from its Ca and Mg contents. The result is the milliliters of your acid which you should apply to each 100 gallons of your water. The factor is calculated to neutralize 80% of the bases in the water. There are 29.6 ml in one U.S. fluid ounce. Divide the number of ml by 29.6 to convert to U.S. fluid ounces.


$$80\% \times \frac{\text{me base}}{\text{L water}} \times \frac{378 \text{ L}}{\text{to be neutralized}} \times \frac{1}{34.7 \text{ N acid}} = 8.7 \times \frac{\text{me base}}{\text{L water}}$$

NOTE: When calculating your rates for larger volumes, be careful not to round off too soon when making conversions.

12. Why not neutralize 100 percent of the bases?

Some of the reasons for not attempting to neutralize 100% of the bases are:

It is not necessary to neutralize all of the bases in order to reduce the problem to insignificant levels. Not trying for 100% neutralization allows some room for error in acid application rates, variability in water, etc. The risk of over-acidifying is not worth the benefit of neutralizing the last 10 or 20 % of the bases. It is poor management to spend money and effort creating new problems by over-reacting to the initial problem.



13. In what kind of irrigation system can I practically inject acid?

Neutralization is relatively easy to accomplish in microirrigation systems. The system must allow careful addition of known volumes of acid to known volumes of water. Since acids can be quite corrosive to metals, the system must be able to withstand this possible adverse effect.

NOTE: It is illegal to inject any chemicals into irrigation systems without appropriate safety devices which will automatically prevent the backflow of water and chemicals to the water supply. This is done to protect our water resources.



14. What kind of acid can I use?

The most commonly used acids are sulfuric, hydrochloric, and phosphoric acid. Other acids could be used but cost and availability usually limit the choices to these three. Phosphoric and sulfuric acids may have some nutritional value but this should be a minor consideration in choosing an acid for water neutralization.

15. What are the dangers of using acids for water neutralization?

Hydrochloric, sulfuric and phosphoric acids are highly toxic materials irritating to the skin, eyes, nose, throat, lungs, and digestive tract. Always wear goggles and chemical resistant (rubber, neoprene, vinyl, etc.) gloves, apron and boots whenever handling these acids. **Acid must be poured into water, never vice versa, and should be done in a well-ventilated area.**

Should a spill or splash occur, remove all clothing and shower immediately. Immediately irrigate eyes with large quantities of water. Seek immediate medical attention.

It is generally advisable to dilute concentrated acid in a nonmetal mixing tank prior to injection into the irrigation system, rather than injecting concentrated acid directly. Most metal fittings, tanks, and other parts of the irrigation system will be damaged by acid, so proper precautions must be taken. Flushing the system after application is frequently sufficient to avoid significant damage. In addition to the dangers involved with handling strong acids there is also the danger of over-application of acid. Excess acid addition could result in injury to tree parts, which come in direct contact with the water, such as leaves. Also, an excessive acidification of the soil could result in tree injury or death. These problems can be avoided by (1) determining the proper amount of acid to apply and (2) monitoring the irrigation system to ensure that the correct amount is applied.

16. How can I assure that I'm adding the correct amount of acid to my water?

Monitoring the pH of the acid-treated water is one way of checking on a daily operational basis. You can do this with a pH meter. Add acid to bring the water pH to between 4.5 and 5.0. Because the neutralization reaction continues slowly over a period of a day or two, the measured pH of the water immediately after acid addition will usually be lower than that measured once the reaction is complete. For monitoring purposes during acid additions, use the pH measured immediately after acid addition as a guide to avoid over-acidifying.

If the pH after treatment is very different from that calculated from the chemical analysis, you may want to have another water sample analyzed.

Summary

1. Have your irrigation water tested.
2. Select an acid of known strength.
3. Determine how much of your acid is needed to neutralize 80% of the bases in your water.
4. Add the calculated amount of acid to your water.
5. Measure the pH of the water as it comes out of the irrigation line.
6. If the pH is not between 4.5 and 5.0, increase or decrease the amount of acid.
7. If the amount of adjustment in Step 6 is more than 15 to 20% of the calculated value, consult a specialist before extended use of the system.
8. Retest the well water and irrigated soil about once a year and keep a record of the test results.





FREQUENTLY ASKED QUESTIONS ABOUT BIOSOLIDS

1) What are Biosolids?

Biosolids are the nutrient-rich solid organic matter recovered from the treatment of domestic sewage in a wastewater treatment facility. Biosolids are a beneficial resource, containing essential plant nutrient and organic matter and are recycled as a fertilizer and soil amendment. When treated and processed, these residuals can be recycled and applied as fertilizer to improve and maintain productive soils and stimulate plant growth.

2) What is the difference between biosolids and sewage sludge?

Sludge is generally used before applicable beneficial recycling criteria have been achieved which normally occurs at the outlet of the stabilization process. It should be used in tandem with a specific process descriptor (e.g., *primary sludge, waste activated sludge, secondary sludge, etc.*)

Biosolids is generally used after applicable beneficial recycling criteria have been achieved, i.e., at the outlet of the stabilization process. Common stabilization processes include the following: aerobic digestion, autothermal thermophilic aerobic digestion (ATAD), anaerobic digestion, composting, alkaline stabilization, thermal drying, including flash, rotary, fluid bed, paddle, hollow-flight, disc, and infrared dryers, thermophilic

pozzolanic fixation, acid oxidation/disinfection, and heat treatment/acid digestion.

3) Why do we have biosolids?

We have biosolids as a result of treating sewage sludge (i.e., the solids generated during the treatment of domestic sewage in a treatment plant) to meet the land application regulatory requirements). Wastewater treatment technology has made our water safer for recreation and seafood harvesting. Forty years ago, thousands of American cities dumped their raw sewage directly into the nation's rivers, lakes, and bays. Through regulation of this dumping, local government now required to treat domestic sewage and to make the decision whether to recycle the solids generated as fertilizer, incinerate them or bury them in a landfill. If the solids meet the regulatory requirements for land application and are recycled, they are biosolids.



4) How are biosolids generated and processed?

Biosolids are generated during the treatment of domestic sewage when

treated further to meet regulatory requirements. The wastewater treatment can actually begin before the wastewater reaches the treatment plant. In many larger wastewater treatment systems, pre-treatment regulations require that industrial facilities pre-treat their wastewater to remove many hazardous contaminants before it is sent to a wastewater treatment plant.

Wastewater treatment facilities monitor incoming wastewater streams to ensure their recyclability and compatibility with the treatment plant process.

Sewage sludge is not generated until domestic sewage is treated in a treatment works, and biosolids are not produced until the sewage sludge meets the land application Part 503 requirements. For these reasons, the treatment of biosolids cannot occur before the domestic sewage reaches the wastewater treatment plant.

Once the wastewater reaches the plant, domestic sewage goes through physical, chemical and biological processes that clean the domestic sewage and remove the solids. If necessary, some of the solids are then treated with lime to raise the pH level to eliminate objectionable odors.

Pathogen reduction (disease-causing organisms, such as bacteria, viruses and parasites) and other organisms capable of transporting disease for the solids usually occur in a different process (e.g., a digester).

5) How are biosolids used?

After treatment and processing, biosolids can be recycled and applied as fertilizer to improve and maintain productive soils and stimulate plant growth. The controlled land application of biosolids completes a natural cycle

in the environment. By treating sewage sludge, it becomes biosolids that can be used as valuable fertilizer, instead of taking up space in a landfill or other disposal facility.

6) Are biosolids safe?

Decades of studies have demonstrated that biosolids can be safely used on food crops. The National Academy of Sciences has reviewed current practices, public health concerns and regulatory standards, and has concluded that "the use of these materials in the production of crops for human consumption when practiced in accordance with existing federal guidelines and regulations, presents negligible risk to the consumer, to crop production and to the environment." In addition, an epidemiological study of the health of farm families using biosolids showed that the use of biosolids was safe.

7) Do biosolids smell?

Biosolids may have their own distinctive odor depending on the type of treatment it has been through. Some biosolids may have only a slight musty, ammonia odor. Others have a stronger odor that may be offensive to some people. Compounds that contain sulfur and ammonia, which are both plant nutrients, cause most odors.

8) Are there regulations for the land application of biosolids?

The federal biosolids rule is contained in 40 CFR Part 503. Biosolids that are to be land applied must meet these strict regulations and quality standards. The Part 503 rule governing the use and disposal of biosolids contains general requirements, numerical limits for metals in biosolids, pathogen and vector attraction reduction standards,

management practices and frequency of monitoring, record keeping and reporting requirements for land applied biosolids as well as similar requirements for sewage sludge that is surface disposed or incinerated. Most recently, Part 503 requirements have been proposed to limit the concentration of dioxin and dioxin like compounds in biosolids to ensure safe land application. Biosolids are one of the most studied materials that have ever been regulated by EPA.

9) Where can I find out more about the regulations?

The biosolids rule is described in the EPA publication, A Plain English Guide to the EPA Part 503 Biosolids Rule. This guide states and interprets the Part 503 rule for the general reader. This guide is also available in hard copy. In addition to the Plain English Guide, EPA has prepared A Guide to the Biosolids Risk Assessments for the EPA Part 503 Rule which shows the many steps followed to develop the scientifically defensible, safe set of rules (also available from EPA in hard copy.)

The cited references provide valuable information about the Part 503 land application requirements. However, if the information in the references is different from the requirements in the Part 503 rule, the Part 503 rule requirements apply. A number of relevant biosolids publications are located on the National Biosolids Partnership's web page at: <http://www.biosolids.org>.

10) How are biosolids used for agriculture?



Biosolids are used to fertilize fields on which crops are grown. Agricultural uses of biosolids that meet strict quality criteria and application rates have been shown to produce significant improvements in crop growth and yield. Nutrients found in biosolids, such as nitrogen, phosphorus and potassium and trace elements such as calcium, copper, iron, magnesium, manganese, sulfur and zinc, are necessary for crop production and growth. The use of biosolids reduces the farmer's production costs and replenishes the organic matter that has been depleted over time. The organic matter improves soil structure by increasing the soil's ability to absorb and store moisture.

Crops use the organic nitrogen and phosphorous found in biosolids very efficiently because these plant nutrients are released slowly throughout the growing season. This enables the crop to absorb these nutrients as the crop grows. This efficiency lessens the likelihood of groundwater pollution of nitrogen and phosphorous.

11) Can biosolids be used for composting?

Yes, biosolids may be composted and sold or distributed for use on lawns and home gardens. Biosolids composted with sawdust, wood chips, yard clippings, or crop residues make excellent mulches and topsoils for horticultural and landscaping purposes. Even after composting, the sewage sludge has to meet the appropriate Part 503 requirements for it to become biosolids that can be applied to lawns and home gardens. Many professional landscapers use composted biosolids for landscaping new homes and businesses. Home gardeners also find composted biosolids to be an excellent addition to planting beds and gardens. Most biosolids compost, are highly desirable products that are easy to store, transport and use.

12) Are there rules about where biosolids can be applied?

To determine whether biosolids can be applied to a particular farm site, a good management practice includes an evaluation of the site's suitability and is generally performed by the land applier. The evaluation examines water supplies, soil characteristics, slopes, vegetation, crop needs and the distances to surface and groundwater.

There are different rules for different classes of biosolids. Class A biosolids contain no detectable levels of pathogens and must meet strict vector attraction reduction requirements and low levels metals contents. The biosolids preparer usually applies for a permit and only have to apply for permits to ensure that these very tough standards have been met. However, the Part 503 requirements

have to be met even if there is no permit. Class B biosolids are treated but still contain detectable levels of pathogens. There are buffer requirements, public access, and crop harvesting restrictions for Class B biosolids. (The land application site restrictions have to be met in all cases where Class B biosolids are land-applied.)

Nutrient management planning ensures that the appropriate quantity of biosolids is land-applied. The biosolids application is specifically calculated to match the nutrient uptake requirements of the particular crop. Nutrient management technicians work with the farm community to assure proper land application and nutrient control.

13) Is EPA pushing the use of biosolids as a fertilizer? Is the federal policy for biosolids driven by economics of disposal?

As a result of its decade-long assessment of biosolids, EPA concluded that recycling biosolids to land was an environmentally responsible solution, when used in accordance with the Part 503 rule. The Federal policies supporting and promoting the beneficial recycling of biosolids are based upon sound science that has demonstrated the benefits of such recycling. These policies are not driven by economics, and the choice of to recycle biosolids remains a local decision.

14) How do the risks associated with biosolids compare with other soil amendments used in agriculture?

A Water Environment Research Foundation (WERF) study completed in 2002 finds that the risks associated with biosolids are no greater than risks

associated with other soil amendments used in agriculture. The project, "Evaluate Risks and Benefits of Soil Amendments Used in Agriculture" (project no. 99-PUM-1), examined the risks and benefits, advantages and potential disadvantages associated with the use of a variety of soil amendments in comparison to chemical fertilizers. Project results indicate that the relative risk to the environment from amendments and fertilizers varies by parameter and shows that known risks from each of the materials studied can be managed. Moreover, these manageable risks must be carefully weighed against the considerable benefits provided by the land application of amendments and fertilizers.

15) Is recycling much cheaper than disposal?

In areas where disposal costs have increased due to shrinking landfill space and increased costs to maintain and monitor landfills, some cities and towns find that recycling biosolids is less expensive than land filling. However, in most cases, land filling is competitive or less expensive than

land application. In such cases, many U.S. communities have made a positive environmental decision to commit to recycling biosolids despite the additional cost. This is especially true where communities have committed to the additional costs of composting or heat drying and pathogen reduction processes for biosolids prior to utilization.

16) Are Biosolids good for the environment?

Recycling biosolids is good for the environment. Organic matter has been recycled for centuries to improve soil fertility and productivity. When properly applied and managed, biosolids can: provide essential plant nutrients; improve soil structure and tilth; add organic matter; enhance moisture retention; and reduce soil erosion.

Biosolids recycling is regulated and encouraged by the United States Environmental Protection Agency and state and local authorities. Research and years of recycling experience have demonstrated that properly managed land application of biosolids is environmentally safe.

- Use of compost, animal manures, biosolids, organic fertilizers, and mulch in agriculture is useful and beneficial.
- Humus, which is the end product of broken down (decayed) organic matter, is an important component of healthy soils and has many great benefits.

Humus improves soil structure, has exceptionally high water-holding capacity and high nutrient storage capacity, supplies plant nutrients, allows more oxygen to enter the soil increasing root growth, increases water penetration into the soil and root development, increases and maintains healthy populations of beneficial micro-organisms and can be very effective at preventing and suppressing plant diseases.

CITRUS RUST MITES



The citrus rust mite (CRM) is an important pest of fruit grown for the fresh market. On some specialty varieties (such as Sunburst tangerine), damage may be particularly severe on stems and foliage, causing leaf injury and possible abscission. Fruit damage is the main concern with other varieties. CRM feeds on green stems, leaves, and fruit. Egg deposition begins within 2 days after the female reaches sexual maturity and continues throughout her life of 14 to 20 days. The female lays one to two spherical transparent eggs per day and as many as 30 during her lifetime. Eggs hatch in about 3 days at 81°F. The newly hatched larva resembles the adult, changing in color from clear to lemon yellow (CRM). After about 2 days at 81°F, molting occurs. The first nymphal stage resembles the larval and requires about 2 days to molt to an adult at the above temperature. The CRM adult has an elongated, wedge-shaped body about three times longer (0.15 mm) than wide. CRM usually is straw to yellow in color. CRM population densities increase in May-July and then decline in late August, but can increase again in late October or early November. Mite densities in the fall rarely approach those early in the summer. During the summer, CRM are more abundant on fruit

and foliage on the outer margins of the tree canopy. Generally, the north bottom of the tree canopy is preferred and supports the highest mite populations. The least favorable conditions for CRM increase are found in the south top of the tree canopy.

Visible characteristics of injury differ according to variety and fruit maturity. When rust mite injury occurs on fruit during exponential growth, before fruit maturity (April to September), epidermal cells are destroyed resulting in smaller fruit. Early season rust mite injury is called "russetting." Rust mite injury to mature fruit (after September) differs significantly from early "russetting." Unlike "russetting" on fruit, fall damaged fruit will polish since the natural cuticle and wax layer remain intact. This condition is known as "bronzing." While the primary effect of fruit damage caused by rust mites appears to be a reduction in grade, other conditions have been associated with severe fruit injury that include reduced size, increased water loss, and increased drop.

Leaf injury caused by feeding of CRM exhibits many symptoms on the upper or lower leaf epidermis. When injury is severe, the upper cuticle can lose its glossy character, taking on a dull, bronze-like color, and/or exhibit patchy yellowish cells in areas of "russetting" that have been degreened by ethylene release during the wounding process. Lower leaf surfaces often show "mesophyll collapse" appearing first as yellow degreened patches (collapsed spongy mesophyll cells) and later as necrotic spots. With the exception of upper leaf epidermal injury to some specialty varieties, such as Ambersweet, Fallglo, and Sunburst, defoliation caused by CRM is rarely severe.

The need for chemical treatments to control rust mites is dictated by numerous biological attributes of the mites, marketing objectives for the fruit, and horticultural practices. These key biological factors include: 1) inherent ability of mites to quickly increase to injurious densities on fruit and sustain the potential for reproductive increase over time; and 2) small size, which makes it difficult to monitor

population densities in the field and detect injurious levels until visible injury has occurred on the fruit. The marketing objective for fruit is particularly important. Cosmetic appearance is a priority for fruit grown for the fresh market. Fruit growth and abscission are not affected until 50% to 75% of the surface has been injured. Thus, there is reduced justification for chemical control of rust mites on fruit grown for processing. Citrus groves producing fruit designated for the fresh market may receive three or four miticides per year, typically during April, June, August, and October. In contrast, groves producing fruit designated for processing receive zero to two treatments per year. Miticides applied for the control of rust mites on fresh fruit varieties are often combined with compatible fungicides in the spring and summer. An alternative approach is using FC 435-66, FC 455-88, or 470 petroleum oil as a fungicide for greasy spot control and to suppress pest mites. From a horticultural perspective, canopy density has an effect on rust mite populations and their ability to increase over a short period of time. The denser the canopy, the less favorable conditions are for a rapid rust mite increase. Since most registered miticides have no ovicidal activity and short residual activity on fruit and foliage, residual control is generally better if the miticide is applied when rust mite adult and egg population densities are low for fresh market varieties. Since external blemishes caused by rust mites, fungal diseases, and wind are less important when fruit are grown for processing, the chemical control strategy for rust mites can be modified significantly. A summer spray is often required for greasy spot control. Use of petroleum oil in place of copper will reduce the likelihood of requiring a subsequent miticide treatment. Further miticide treatment may be unnecessary. However, a second petroleum oil application may be required for greasy spot control on summer flush. Many scientific methods for sampling or scouting rust mite populations have been described. Of these, three general approaches are in widespread use: 1) determining the percentage of fruit and/or

leaves infested with rust mites, 2) qualitative rating scales and 3) individual adult mite counts taken from fruit on randomly selected trees. These sampling approaches are similar in that they are designed to avoid bias by randomly selecting different representative areas within a grove for sampling, avoiding border rows, and selecting fruit and/or leaves within a tree randomly.

One sampling method based on rust mite density (rust mites/square centimeter [cm^2]) is described.

Processed Fruit: Initiate rust mite monitoring in April on leaves and fruit through casual observations and continue every 2 to 3 weeks throughout the fruit season. Select trees at random and within uniformly distributed areas throughout a 10- to 40-acre block representing a single variety with uniform horticultural practices. Avoid sampling adjacent trees. Fruit should be sampled at random representing the four quadrants of the tree and taken midway in the canopy (between interior and exterior). One fruit surface area should be examined midway between the sun and shade areas. The number of rust mites per cm^2 should be recorded and averaged for the 10 acres, represented by 20 trees with four fruit per tree or 80 readings per 10 acres. Six rust mites/ cm^2 would be a planning threshold where pesticide intervention may be required within 10 to 14 days. Ten rust mites/ cm^2 would be an action threshold where treatment would be required as soon as possible.

Fresh Fruit: Similar to above except monitor every 10 to 14 days with an average of 2 CRM/ cm^2 as an action threshold.

For more information, go to:

<http://www.crec.ifas.ufl.edu/extension/pest/PDF/2016/Rust%20Mites.pdf>

TABLE 1. CONTROL THRESHOLDS AND APPROPRIATE SAMPLE SIZES FOR 10 ACRES

If the control threshold is:	Sample size (Sample trees should be uniformly scattered across a 10-acre block. Do not sample adjacent trees.)
5 mites/leaf	Examine 4 leaves/tree from 6 trees/area from 4 areas/10 acres = 96 leaves on 24 trees/10 acres
8 mites/leaf	Examine 4 leaves/tree from 6 trees/area from 3 areas/10 acres = 72 leaves on 18 trees/10 acres
10 mites/leaf	Examine 4 leaves/tree from 5 trees/area from 2 areas/10 acres = 40 leaves on 10 trees/10 acres
15 mites/leaf	Examine 4 leaves/tree from 4 trees/area from 2 areas/10 acres = 32 leaves on 8 trees/10 acres

TABLE 2. CITRUS MITICIDE SELECTION*

Supplemental (early Spring)	Post Bloom	Summer	Fall	Supplemental Fall
--	--	Agri-mek + oil	--	--
--	--	--	Comite	Comite
Envidor	Envidor	Envidor	Envidor	Envidor
--	Petroleum oil	Petroleum oil	Petroleum oil	--
--	--	--	Sulfur	Sulfur
--	--	Micromite	Micromite	--
--	--	--	Nexter	Nexter
Movento	Movento	Movento	--	--
Vendex	Vendex	--	Vendex	Vendex

*Except for petroleum oil, do not use the same miticide chemistry more than once a year.

GREASY SPOT FUNGAL DISEASE

Management of greasy spot must be considered in groves intended for processing and fresh market fruit. Greasy spot is usually more severe on leaves of grapefruit, pineapples, Hamlins, and tangelos than on Valencias, Temples, Murcotts, and most tangerines and their hybrids.

Greasy spot spores germinate on the underside of the leaves and the fungus penetrates through the stomates (natural openings on lower leaf surface). Warm humid nights and high rainfall, typical of Florida summers, favor infection and disease development.



On processing Valencias, a single spray of oil (5-10 gal/acre) or copper + oil (5 gal/acre) should provide acceptable control when applied from mid-May to June. With average quality copper products, 2 lb of metallic copper per acre usually provide adequate control. The strobilurin fungicides (Abound, Gem, Headline or Quadris), as well as Enable 2F, are also suitable with or without petroleum oil. On early and mid-season oranges and grapefruit for processing, two sprays may be needed especially in the southern part of the state where summer flushes constitute a large portion of the foliage. Two applications also may be needed where severe defoliation from greasy spot occurred in the previous year. In those cases, the first spray should be applied from mid-May to June and the second soon after the major summer flush has expanded. Copper fungicides provide a high degree of control more consistently than oil sprays. Control of greasy spot on late summer flushes is less important than on the spring and early summer growth flushes since the disease develops slowly and defoliation will not occur until after the next year's spring flush. Thorough coverage of the underside of leaves is necessary for maximum control of greasy spot, and higher spray volumes and slower tractor speeds may be needed than for control of other pests and diseases.



The program is essentially the same for fresh fruit. That is, a fungicide application in May-June and a second in July should provide control of rind blotch.

A third application in August may be needed if rind blotch has been severe in the grove. Petroleum oil alone is less effective than other fungicides for control of greasy spot rind blotch (GSRB). Heavier oils (455 or 470) are more effective for rind blotch control than are lighter oils (435).

Copper fungicides are effective for control of GSRB, but may result in fruit spotting especially if applied at high rates in hot, dry weather or if applied with petroleum oil. If copper fungicides are applied in summer, they should be applied when temperatures are moderate, at rates no more than 2 lb of metallic copper per acre, without petroleum oil or other additives, and using spray volumes of at least 125 gal/acre. Enable 2F can be applied for greasy spot control at any time but is especially indicated in mid to late summer for rind blotch control.

The strobilurin fungicides (Abound, Gem, Headline, Pristine or Quadris Top) or Enable 2F can be applied at any time to all citrus and provide effective control of the disease on leaves or fruit. Use of a strobilurin (Abound, Gem, Headline, Pristine or Quadris Top) is especially indicated in late May and early June since it will control both melanose and greasy spot and avoids potential fruit damage from the copper fungicides at that time of year. A strobilurin fungicide should not be applied

more than once a year for greasy spot control. Addition of petroleum oil increases the efficacy of these products.

•Processed fruit

May-June

- Petroleum oil (455, 470) 5-10 gal
- Cu fungicides 2-4 lb metal
- Abound, Gem, Headline + 5 gal oil
- Pristine
- Quadris Top
- Enable

July

- Petroleum oil (455, 470) 5-10 gal
- Cu fungicides 2-4 lb metal
- Abound, Gem, Headline + 5 gal oil
- Pristine
- Quadris Top
- Enable

•Fresh fruit

May-June

- Petroleum oil (455, 470) 10 gal
- Cu fungicides < 2 lb metal, No oil
- Abound, Gem, Headline + 5 gal oil
- Pristine
- Quadris Top
- Enable

July

- Petroleum oil (455, 470) 10 gal
- Cu fungicides < 2 lb metal
- Abound, Gem, Headline + 5 gal oil
- Pristine
- Quadris Top
- Enable 8 oz

For more information on greasy spot, go to:

<http://www.crec.ifas.ufl.edu/extension/pest/PDF/2016/Greasy%20Spot.pdf>

Flatwoods Citrus

If you did not receive the *Flatwoods Citrus* newsletter and would like to be on our mailing list, please check this box and complete the information requested below.

If you wish to be removed from our mailing list, please check this box and complete the information requested below.

Please send: Dr. Mongi Zekri
Multi-County Citrus Agent
Hendry County Extension Office
P.O. Box 68
LaBelle, FL 33975

Subscriber's Name: _____

Company: _____

Address: _____

City: _____ State: _____ Zip: _____

Phone: _____

Fax: _____

E-mail: _____

Racial-Ethnic Background

American Indian or native Alaskan

Asian American

Hispanic

White, non-Hispanic

Black, non-Hispanic

Gender

Female

Male