

Hendry County Extension, P.O. Box 68, LaBelle, FL 33975 (863) 674 4092

# Flatwoods Citrus



**Vol. 25, No. 7**

**July 2022**

**Dr. Mongi Zekri**  
**Multi-County Citrus Agent, SW Florida**



## ***Table of Contents***

July 2022 Citrus Seminar	2
CEUs for Pesticide License Renewal	3
Flatwoods Citrus Newsletter Sponsors – Thank you!	4-6
El Niño/Southern Oscillation (ENSO) Diagnostic Discussion	7
Flooding Injury	8-9
Water Table Measurement and Monitoring	10-11
Resetting in Citrus Groves	12-15
Danger of Heat Stress	16-17
Managing Heat Stress	18-19
Pesticide Resistance and Resistance Management	20-21
Citrus Spray Programs	22

**Please mark your calendar and plan to attend**

## **July 2022 Citrus Seminar** via Zoom only

The Zoom link <https://ufl.zoom.us/j/91579830879?pwd=NitXL2toOHAXSWVMcHM2b0lVTmIKUT09>

After registering, you will receive a confirmation email containing information about joining the Zoom meeting.

No registration fee

Date & Time: **Wednesday, July 20, 2022, 9:00 AM – 10:00 AM**

Title: **Citrus Huanglongbing is an immune-mediated plant disease and its implications in HLB management**

Speaker: **Dr. Nian Wang**, Professor in microbiology and cell science, UF/IFAS Citrus Research & Education Center, Lake Alfred

Our recent study demonstrate that Citrus Huanglongbing is a pathogen-triggered immune disease. We discovered that CLAs infection of citrus stimulates systemic and chronic immune response in phloem tissues including reactive oxygen species (ROS) production. Systemic cell death of phloem tissues is caused by excessive and chronic ROS production triggered by CLAs. Consequently, cell death of phloem tissues causes HLB symptoms. The finding of citrus HLB as an immune-mediated plant disease helps guide the battle against this notorious disease. It seems likely that horticultural and genetic approaches that suppress ROS damages can manage HLB. I will talk about the shove-ready approaches including inducing the activities of antioxidant enzymes via application of micronutrients (B, Fe, Mo, Ni, and Zn), promoting plant growth using plant growth hormones, such as gibberellin, and suppressing reactive oxygen species (ROS) damages using antioxidants, such as uric acid which is yet to be labeled on citrus. I will talk about our current progress on generating HLB resistant/tolerant citrus varieties. I will also talk about other experiments that we have conducted that might be interested to citrus industry including trunk injection with antimicrobials.

Coordinator: Dr. Mongi Zekri, Multi-County Citrus Extension Agent, UF-IFAS

1 CEU for pesticide license renewal

1 CEU for certified crop advisors

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Employment Opportunity - Affirmative Action Employer authorized to provide research, educational information and other services only to individuals and institutions that function without regard to race, color, sex, age, handicap or national origin.

U.S. DEPARTMENT OF AGRICULTURE, COOPERATIVE EXTENSION SERVICE, UNIVERSITY OF FLORIDA, IFAS, FLORIDA A. & M. UNIVERSITY COOPERATIVE EXTENSION PROGRAM, AND BOARDS OF COUNTY COMMISSIONERS COOPERATING.



## CEUs for pesticide license renewal

Earn CEU Credits NOW online through Southeast AgNet & Citrus Industry magazine

<http://citrusindustry.net/ceu/>

The following series of articles and quizzes are available with their expiration dates noted:

- #2 **How Weather Affects Pesticide Applications** (4/30/23)
- #1: **Increasing Pesticide Effectiveness With Adjuvants** (1/31/23)
- #4: **Protecting People From Pesticide Exposure** (10/31/22)
- #3: **Before You Spray** (7/31/22)

***Each article grants one General Standards (Core) CEU when submitted and approved toward the renewal of a Florida Department of Agriculture and Consumer Services restricted-use pesticide license.***

FYI, there are also CORE CEU available at Growing Produce

<http://www.growingproduce.com/crop-protection/ceu-series/>

<http://www.growingproduce.com/crop-protection/ceu-series/>

Online Pesticide CEUs

<https://pested.ifas.ufl.edu/ceu/>



**Special Thanks to sponsors of the "Flatwoods Citrus" newsletter for their generous contribution and support. If you would like to be among them, please contact me at 863 674 4092 or maz@ufl.edu**



**Sam Thayer**  
President

P.O. Box 1849  
Dundee, FL 33838  
Phone: (863) 439-3667  
Fax: (863) 439-6608

[www.maxijet.com](http://www.maxijet.com)  
[sam@maxijet.com](mailto:sam@maxijet.com)

The Standard of Quality In Low-Volume Irrigation



**Frank Miele**

**Office: 863 357 0400**

**Cell: 954 275 1830**

**Fax: 863 357 1083**

**E-mail: [famiele1@aol.com](mailto:famiele1@aol.com)**



**Brad Lang**

**Territory Sales Manager**

**Phone: 229 894 0568**

**[brad.lang@nufarm.com](mailto:brad.lang@nufarm.com)**

**AGRICULTURAL  
LAND BROKERAGE**

**LSI  
COMPANIES**

Call our Ag specialists, Billy Rollins & Hunter Ward

**[LSIcompanies.com](http://LSIcompanies.com) | 239.489.4066**

**Clint Wise Jr.**

**AGLIME SALES, INC.**

P.O. Box 60

Babson Park, FL 33827

**863-241-9007**

**[clint.wise@aglimesales.com](mailto:clint.wise@aglimesales.com)**

**MIKE HERRINGTON**  
863-412-2067  
[MikeH@amvac.com](mailto:MikeH@amvac.com)

CITRUS FIX™ • Krovar® DF • DIBROM®  
B Sure® • DEADLINE® • ECOZIN®

**Justin Newsome**

**syngenta**

**Mobile: 863 221 9681**

**[justin.newsome@syngenta.com](mailto:justin.newsome@syngenta.com)**

**[www.syngenta.com](http://www.syngenta.com)**



**AGRICULTURAL**  
RISK MANAGEMENT LLC

*Growing with you!*

**Trevor Troyer**  
Office: 239 789 4742  
Cell: 239 810 0138  
E-mail: [ttroyer@agriskmgmt.com](mailto:ttroyer@agriskmgmt.com)  
[www.agriskmgmt.com](http://www.agriskmgmt.com)



**“MAKE CITRUS GREAT AGAIN”™**

**863-439-2877**  
[www.TheTreeDefender.com](http://www.TheTreeDefender.com)

PATENT PENDING

**Special Thanks to sponsors of the “Flatwoods Citrus” newsletter for their generous contribution and support. If you would like to be among them, please contact me at 863 674 4092 or [maz@ufl.edu](mailto:maz@ufl.edu)**



**BLACKSMITH**  
BIOSCIENCE

**Jack Kilgore**  
Technical Sales Rep, SE US

Office: 7150 East Brentwood Road Ft. Myers, FL 33919 (239) 707-7677  
[g8trmanjek@comcast.net](mailto:g8trmanjek@comcast.net)  
[www.blacksmithbio.com](http://www.blacksmithbio.com)

**TIGER-SUL IS A PROUD SPONSOR OF FL CITRUS GREENING RESEARCH**

Discover how TIGER Greening Guard Citrus Mix is helping maintain strong returns on investments and keeping HLB infected trees as healthy and productive as possible, for as long as possible.

**Kenneth Waters**  
Tiger-Sul Products, LLC  
[kwaters@tigersul.com](mailto:kwaters@tigersul.com)  
Phone: 850.501.6127  
[www.tigersul.com](http://www.tigersul.com)



**ASK FOR TIGER!**

**Heath Prescott**

**KeyPlex®**

Toll Free: 800 433 7117  
Mobile: 863 781 9096  
[heath@keyplex.com](mailto:heath@keyplex.com)

**NICHINO AMERICA**  
**Scott Croxton**  
[scroxton@nichino.net](mailto:scroxton@nichino.net)  
**Samuel S. Monroe**  
[smonroe@nichino.net](mailto:smonroe@nichino.net)  
[www.nichino.net](http://www.nichino.net)



**CENTRAL**  
LIFE SCIENCES

**Extinguish**  
PLUS

**EXTINGUISH**  
Professional Fire Ant Bait

[www.CentralAntControl.com](http://www.CentralAntControl.com)  
Rodney Morrow  
(352) 651-2780  
[rmorrow@central.com](mailto:rmorrow@central.com)




**Plant Food Systems, Inc.**  
**P.O. Box 775**  
**Zellwood, FL 32798**  
**Tel: 407 889 7755**

POLYON® Controlled-Release fertilizer makes you more efficient. *Ask me how!*

POWERED BY  
**POLYON**  
 Controlled-Release Fertilizer

**Trey Whitehurst (863)-633-8711**  
[www.harrells.com](http://www.harrells.com)



**AMERI COP<sup>40 DF</sup>**

**COPPER HYDROXIDE**  
**FUNGICIDE AND BACTERICIDE**

Jack Kilgore • g8trmanjek@comcast.net • (239)-707-7677  
[www.iqvagro.com](http://www.iqvagro.com)

OMRI LISTED

NOT ALL LIQUID COPPERS ARE CREATED EQUAL

**Badge<sup>SC</sup>**  
 Fungicide/Bactericide

**BART HOOPINGARNER**  
**941.737.7444**  
[BHOOPINGARNER@GOWANCO.COM](mailto:BHOOPINGARNER@GOWANCO.COM)

Badge<sup>SC</sup> is a registered trademark of Isagro USA, Inc.  
 Always read and follow label directions.

**Gowan**  
 USA  
[GOWANCO.COM](http://GOWANCO.COM)

**FMC Corporation**  
**Ed Early, 239-994-8594**  
[Edward.Early@fmc.com](mailto:Edward.Early@fmc.com)  
**Daren Sapp, 863 840 4600**  
[Daren.sapp@fmc.com](mailto:Daren.sapp@fmc.com)  
**Brent Johnson, 941 243 3379**  
[Dennis.Johnson@fmc.com](mailto:Dennis.Johnson@fmc.com)



**Timac AGRO**  
 USA  
 THINK DIFFERENT... FARM BETTER

**CHARLES McCARTNEY**  
 SOUTHEAST BUSINESS UNIT DIRECTOR

[CMCCARTNEY@TIMACUSA.COM](mailto:CMCCARTNEY@TIMACUSA.COM)  
 941-219-9063  
[WWW.US.TIMACAGRO.COM](http://WWW.US.TIMACAGRO.COM)

# EL NIÑO/SOUTHERN OSCILLATION (ENSO) DIAGNOSTIC DISCUSSION

issued by

**CLIMATE PREDICTION CENTER/NCEP/NWS**  
**and the International Research Institute for Climate and Society**  
9 June 2022

## **ENSO Alert System Status: [La Niña Advisory](#)**

**Synopsis: Though La Niña is favored to continue through the end of the year, the odds for La Niña decrease into the Northern Hemisphere late summer (52% chance in July-September 2022) before slightly increasing through the Northern Hemisphere fall and early winter 2022 (58-59% chance).**

During May, below-average sea surface temperatures (SSTs) continued across most of the central and eastern equatorial Pacific Ocean [\[Fig. 1\]](#). However, negative SST anomalies weakened during the past month, as reflected by the Niño indices, which ranged from -0.6°C to -0.9°C during the past week [\[Fig. 2\]](#). Subsurface temperatures anomalies (averaged between 180°-100°W and 0-300m depth) also weakened with values returning to near zero [\[Fig. 3\]](#). Below-average subsurface temperatures persisted near the surface to at least ~75m depth from the central to the eastern equatorial Pacific Ocean, with above-average temperatures continuing at depth (~100 to 200m) in the western and central Pacific Ocean [\[Fig. 4\]](#). Low-level easterly wind anomalies prevailed in the east-central equatorial Pacific, while upper-level westerly wind anomalies continued over most of the equatorial Pacific. Convection was suppressed over the western and central Pacific and was weakly enhanced over parts of Indonesia [\[Fig. 5\]](#). Overall, the coupled ocean-atmosphere system continues to reflect La Niña.

The most recent IRI/CPC plume average for the Niño-3.4 SST index forecasts La Niña to persist into the Northern Hemisphere winter 2022-23 [\[Fig. 6\]](#). This is now in greater agreement with the forecast consensus this month, which also predicts La Niña to continue into the winter. However, it is clear that recent observed oceanic and atmospheric anomalies have weakened and this is anticipated to continue through the summer. Uncertainty remains over whether La Niña may transition to ENSO-neutral during the summer, with forecasters predicting a 52% chance of La Niña and a 46% chance of ENSO-neutral during July-September 2022. After this season, the forecast is for renewed cooling, with La Niña favored during the fall and early winter. In summary, though La Niña is favored to continue through the end of the year, the odds for La Niña decrease into the Northern Hemisphere late summer (52% chance in July-September 2022) before slightly increasing through the Northern Hemisphere fall and early winter 2022 (58-59% chance; click [CPC/IRI consensus forecast](#) for the chances in each 3-month period).

This discussion is a consolidated effort of the National Oceanic and Atmospheric Administration (NOAA), NOAA's National Weather Service, and their funded institutions. Oceanic and atmospheric conditions are updated weekly on the Climate Prediction Center web site ([El Niño/La Niña Current Conditions and Expert Discussions](#)). Additional perspectives and analysis are also available in an [ENSO blog](#). A probabilistic strength forecast is [available here](#). The next ENSO Diagnostics Discussion is scheduled for 14 July 2022.

**Climate Prediction Center**  
**National Centers for Environmental Prediction**  
**NOAA/National Weather Service**

## FLOODING INJURY

Almost all citrus trees grown in southwest Florida are located on high water table, poorly drained soils. Water management on poorly drained soils is difficult and expensive because during heavy rains in the summer, excess water must be removed from the rootzone and in periods of limited rainfall, irrigation is needed. On these soils, drainage is as important as irrigation. The concept of total water management must be practiced. If either system—irrigation or drainage—is not designed, operated, and maintained properly, then the maximum profit potential of a grove cannot be achieved. Both surface and subsoil drainage is necessary to obtain adequate root systems for the trees.

Roots, like the rest of the tree, require oxygen for respiration and growth. Soils in Florida typically contain 20-21 % oxygen. When flooding occurs, the soil oxygen is replaced by water. This condition causes tremendous changes in the types of organisms present in the soil and in the soil chemistry.

Flooding injury would be expected if the root zone were saturated for 3 days or more during extended summer rains at relatively high soil temperatures (86-95° F). Flooding during the cooler December-March period can be tolerated for several weeks at low soil temperatures (< 60° F). The rate of oxygen loss from the soil is much greater at high than at low temperatures. The potential for damage to roots is less obvious but equally serious when the water table is just below the surface. Flooding stress is much less when water is moving than when water is stagnant. The use of observation wells is a very reliable method for evaluating water-saturated zones in sites subject to chronic flooding injury.



Short-term estimates of flooding stress can be obtained by digging into the soil and smelling soil and root samples. Sour odors indicate an oxygen deficient environment. The presence of hydrogen sulfide (a disagreeable rotten egg odor) and sloughing roots indicate that feeder roots are dying. Under flooded conditions, root death is not exclusively associated with oxygen deficiency. Anaerobic bacteria (the kind that can grow only in the absence of oxygen) develop rapidly in flooded soils and contribute to the destruction of citrus roots. Toxic sulfides and nitrites formed by anaerobic sulfate- and nitrate-reducing bacteria are found in poorly drained groves. Sulfate-reducing bacteria require both energy and sulfates in order to change sulfates to sulfides. The best sources of energy have been found to be certain organic acids contained in citrus roots, grass roots, and buried pieces of palmetto. Thus, citrus roots can contribute to their own destruction by being an energy source for these bacteria.



Symptoms of flooding injury may occur within a few days or weeks, but usually show up after the water table has dropped and the roots become stranded in dry soils. Leaf wilting, leaf drop, dieback, and chlorosis patterns may develop and tree death may occur. Trees subjected to chronic flooding damage are stunted with sparse canopies, dull colored, small leaves and produce low yields of small fruit. New flushes of growth will have small, pale leaves due to poor nitrogen uptake by restricted root systems. Usually, the entire grove is not affected, but most likely smaller more defined areas will exhibit the symptoms. Striking differences in tree condition can appear within short distances associated with only slight changes in rooting depths. Water damage may also be recognized by a marked absence of feeder roots and root bark, which is soft and easily sloughed.

With acute water damage, foliage wilts suddenly followed by heavy leaf drop. Trees may totally defoliate and actually die, but more frequently partial defoliation is followed by some recovery. However, such trees remain in a state of decline and are very susceptible to drought when the dry season arrives because of the shallow, restricted, root systems. Moreover, waterlogged soil conditions, besides debilitating the tree, are conducive to the proliferation of soil-borne fungi such as *Phytophthora* root and foot rot. These organisms cause extensive tree death especially in poorly drained soils.

Water damage may usually be distinguished from other types of decline by a study of the history of soil water conditions in the affected areas. Areas showing water damage are usually localized and do not increase in size progressively as do areas of spreading decline. Foot or root rot symptoms include a pronounced chlorosis of the leaf veins caused by root damage and girdling of the trunk. Lesions also appear on the trunk usually near the soil level (foot rot) or roots die and slough-off (root rot). Flood damage does not produce lesions. Trees with blight or CTV are usually randomly distributed within the grove and diagnostic tests are available to distinguish them from water-damaged trees.

Citrus trees respond physiologically to flooding long before morphological symptoms or yield reductions appear. Photosynthesis and transpiration decrease within 24 hours of flooding and remain low as flooding persists. Water uptake is also reduced which eventually translates to decreased shoot growth and yields.

It is both difficult and costly to improve drainage in existing groves, so drainage problems should be eliminated when the grove area is prepared for planting by including a system of ditches, beds and/or tiling. Growers should not depend on the slight and often unpredictable differences in rootstock tolerance to waterlogging to enable trees to perform satisfactorily under such conditions. Trees, irrespective of scion and rootstock cultivars, should be planted under the best drainage conditions possible. Drainage ditches should be kept free of obstruction through a good maintenance program including chemical weed control. Tree recovery from temporary flooding is more likely to occur under good drainage structure maintenance conditions.

Do not disk a grove if trees were injured by flooding. Irrigation amounts should be reduced, but frequencies should be increased to adequately provide water to the depleted, shallow root systems. Soil and root conditions should be evaluated after the flooding has subsided. Potential for fungal invasion should be determined through soil sampling and propagule counts. If there is a *Phytophthora* problem, the use of certain fungicides can improve the situation.

## WATER TABLE MEASUREMENT AND MONITORING



Most flatwoods citrus soils have a restrictive layer that can perch the water table and significantly affect tree water relations. To optimize production and tree health, the level of this water table should be monitored and maintained within an optimal zone. Simple and practical observation wells can normally produce adequate information.

**Water Table Behavior.** The water table under flatwoods citrus may rise rapidly in response to either rainfall or irrigation because sandy soils are highly conductive to water flow. A general rule of thumb is that 1 inch of rain will cause the water table to rise about 10 inches in fine textured soils, 6 inches in most of the flatwoods sandy soils, and 4 inches in coarse sands. It may take 4 to 6 days for the water table to return to its desired levels following rains of 1 inch or more.

**Observation Wells.** A water table observation well is made with a porous casing buried vertically in the ground. It permits the groundwater level to rise and fall inside it as the water level in the adjacent soils. Observation wells with a simple float indicator can provide rapid evaluation of shallow water table depths. The float and indicator level move with the water table, allowing an above-ground indication of the water level. Water table observation wells installed in flatwoods soils usually penetrate only to the depth of the restrictive (argillic or spodic) layer. Typically this layer is within 30 to 48 inches of the soil surface.

**Well Construction.** The basic components of the well itself include a short section of 3-inch perforated PVC pipe (3-5 ft long), 3-inch PVC cap, screening material, a float, indicator rod, and small stopper.

The indicator rod can be a dowel,  $\frac{1}{2}$ -inch PVC pipe (thin wall) or microsprinkler extension stake. Dowels are a poor choice since they require painting and will rot out near the float within a few years. The float is typically a 2½-inch fishing net float or a 500 ml (approximately 2½ in. diameter x 6 in. high) polyethylene bottle with a 28-mm (1.1 in.) screw cap size. The float assembly can be constructed by inserting the microsprinkler extension stake into the fishing float or  $\frac{1}{2}$ -inch pipe into the polyethylene bottle.

The bottle neck provides a snug fit for the stake and no sealant is required. The hole in the cap should be drilled slightly larger than the indicator stake to serve as a guide for the float assembly. Fittings should not be glued so that components can be easily disassembled for cleaning or replacement.

Observation well casings are constructed from 3-in. diameter PVC pipe (Class 160). A circular saw or drill can be used to perforate the pipe prior to installation. Perforations should be staggered in rows around the pipe to allow flow into the well from the sides in addition to the bottom. Perforations totaling about 5% of the well's surface area are adequate for sandy soils

encountered in the flatwoods. No perforations should be made within 12 inches of the surface in order to minimize the chances of ponded water from high intensity storms creating flow channels into perforations near the soil surface.

The pipe should be wrapped (sides and bottom) with a screening material to prevent soil particles from moving into the well. Materials such as cheesecloth, polyester drain fabric, and fiberglass screen have been used successfully as filters. The filter material should be taped in place with duct tape. A 3-inch soil auger can be used to bore holes for the wells. When possible, the observation wells should be installed when no water table is present in order to minimize chances of the well sides sloughing into the bore as it is dug.

When a water table is present, it is easiest to install the well by starting off with a larger diameter pipe. For a 3-inch observation well, a 4-inch installation pipe (Sch 40 preferred) will be needed. The installation pipe should be cut at least 6 inches longer than the intended depth of well. Holes (½-inch diameter) should be drilled in the sides of the pipe opposite each other about 1½ inches from the top of the pipe. These will be used to aid in removing the pipe from the soil after the observation well is installed. Auger a hole in the soil until it begins to slough in (when the water table is reached). The 4-inch pipe should then be forced into the hole. A 3-inch auger can then be used to remove soil from within the 4-inch casing. As soil is removed, the casing needs to be forced downward to keep the hole from sloughing. Continue to remove soil from inside the casing until the appropriate depth is achieved (typically when hardpan material begins to be excavated).

The well casing pipe should be cut to length and installed in the hole so that it extends 2 to 6 inches above the soil surface. Care should be taken to ensure that the casing is installed plumb to minimize binding of the float assembly. If a 4-inch installation pipe was used to excavate the hole, it needs to be removed. A ½-inch rod can be inserted through the holes that were drilled in the top of the 4-inch pipe. If the pipe cannot be removed easily by hand, a chain can be attached to the rod and attached to a high-lift jack. Usually, after jacking the installation pipe up about a foot, the pipe can be easily removed by hand. The soil should be backfilled around the observation well casing and tamped to compact the soil and get a tight fit between the soil and the sides of the pipe.

A measurement should be taken of the distance from the bottom of the well to the soil surface. The float assembly can then be lowered into the well. Make sure that the indicator rod and float do not bind against the sides of the observation well. The well is now ready for calibration.

**Calibration.** A mark on the indicator stake or rod should be made at the top of the well when the float is at the bottom of the well. This level is the reference mark for the well depth. The indicator stake or rod can then be marked with major divisions (feet) and minor divisions (inches) for easy reading of the water table depth. These rings can be painted at appropriate intervals using different colors for major and minor divisions. Marks painted at 2-inch increments provide enough accuracy for most users.

The mark at the upper level is dependent on the depth of the water furrow and root depth. The upper depth should be selected so that water does not pond in water furrows and it should be at least 6 inches below the bottom of the root zone to prevent root pruning. Observations over time will help to determine the water table level depth that will prevent root damage or excessive wetness in the root zone.

# RESETTING IN CITRUS GROVES



For maximum efficiency of a production unit or grove, it is essential that every tree space is occupied by a healthy and productive tree. The average annual tree loss across the Florida citrus industry is currently around 6%. However, the extent of tree loss among individual groves can vary from 2 to 12% or more. Prompt replacement of unproductive trees means higher average long-term returns from the grove. If the declining trees remain in the grove, they keep getting weaker and yield less fruit each year and therefore the potential production capacity for the grove keeps declining even though production costs remain the same or even increase. It is very important to remove and replace such trees once it is clear that they are declining and they are no longer economically profitable. However, the reason for the decline should be determined and the condition should be corrected so that the replacement tree does not suffer the same fate.

Resetting should be considered if the tree is affected by an incurable disease such as blight, tristeza, or citrus greening. The resetting program should be conducted regularly rather than being delayed until serious losses in production have occurred. Resets should be planted with the same cultivar already in the block. Usually, it is more economical to keep resetting and not to push the entire block unless the cultivar and/or the tree spacing between rows is an undesirable one. Replanting in a mature grove seems justified only when a minimum of 8 ft between canopy driplines, (not from trunk-to-trunk), is available for canopy development of the new trees.

Replacement of dead, diseased, and declining trees in Florida citrus groves should always be an important part of the total production program. Today, tree replacement is more important than ever since overhead and production costs are dramatically increasing and a full stand of productive trees is essential to maximize production and profits. Freezes, blight, tristeza, Phytophthora, Diaprepes, and other pests and diseases have been particularly troublesome to Florida citrus growers for the last two decades. Citrus canker and greening have been devastating citrus groves since their introduction to Florida. Extensive tree losses coupled with the economic necessity of regular resetting have caused many growers to investigate ways to achieve new efficiencies in reset management.

## NOT AN EASY TASK

Caring for young citrus trees is always troublesome because they require far more attention than larger, established trees. Florida's sandy soils, high summer temperatures, possible low winter temperatures, and scattered rainfall patterns complicate young tree care by forcing growers to protect, fertilize, and weed young trees regularly or face extensive losses. Young trees are more sensitive and more attractive to pests than mature trees due to high levels of vegetative growth. Therefore, special care is needed to insure pests are adequately controlled. Resets often present an even greater problem because trees are usually scattered throughout a block of larger trees, where they compete with large, full-grown trees for limited supplies of water, nutrients, and sometimes sunlight. Scattered resets frequently have serious weed problems since removal of the previous tree allows the area to receive more sunlight and provides more favorable conditions for weed growth. Since resets are usually scattered throughout a block of much larger trees, they are often difficult to locate and may be accidentally overlooked, resulting in inadequate care. Researchers, growers, and production managers are continually developing and improving methods of dealing with reset care.

## PLANNING THE RESET PROGRAM

Grove managers should include tree removal and resetting as a routine part of the production program and assign special crews to deal specifically with young tree care. Planning ahead is very important because there is often a lag period between the time when replacement trees are ordered and when they are received. The wait time for the most desired rootstock and scion combination may be as great as 1 to 2 years, so replacement tree needs should be anticipated (when possible) and orders placed so they can be obtained when needed.

## PURCHASING TREES

High quality reset trees are essential for maximum young tree growth. These young trees will be placed in an intensely competitive situation and may sometimes receive less than ideal care, so there is no room for compromising tree quality. Only healthy and properly sized trees from registered sources should be purchased since the initial cost is only a small fraction of the total cost of bringing such a tree into production.

## SITE PREPARATION

The planting site should be well prepared. Weeds should be removed before planting. At a minimum, a non-residual herbicide should be applied to the reset area to get weeds under control before the young tree is planted.

Planting sites should be prepared well in advance of receipt of the trees. Ideally, trees should be planted on the same day they are received. Under no circumstances should trees be allowed to dry out. To minimize root desiccation and damage, they should be kept cool and moist until they are planted.

## PLANTING THE TREES

Trees should be removed from the container and inspected for evidence of pot-binding. Make several vertical slashes about one inch deep through the root ball to encourage root branching. These slashes also allow the potting soil and roots to interface more closely with the soil in the planting site. It may be easier to cut some of the roots with pruning shears and pull them so they protrude from the ball.

A common problem with nursery trees is that the potting mixture is often highly organic. Such materials form areas, which are difficult to permeate with water after the young tree is planted in sandy soils and irrigated. The outer third of the organic ball should be removed so that the outer roots are exposed and can extend into the soil in which the tree is planted. Otherwise, the tree may not grow off quickly and satisfactorily.

## WEED CONTROL

Keeping weeds under control during the establishment period of the reset is very important. Weeds compete with young citrus trees for water, nutrients, soil applied pesticides, and sunlight and they must be properly controlled. Weed control around a reset site should be considered at pre-plant, early post-plant, and after the tree is established. Control of weeds prior to planting should be provided. If residual herbicides are used, they should be used at proper rates and at least 30 days in advance of planting so that residues do not impact reset growth. Prior to planting, contact or growth regulating herbicides may be preferred since they do not leave residual effects in the soil.

Weed control during the establishment period or approximately the first year is frequently quite difficult. Hand labor is scarce and expensive. Trunk damage by hoes or other cultivation equipment further compounds the problem. Chemical weed control provides at least a partial solution to the problem during this establishment period. There is now a fairly wide selection of residual herbicides available, which can be used around young trees. These materials should be applied at reduced rates. Be sure to read labels carefully for restrictions on the use of herbicidal materials around young trees.

After the reset has been planted for a year or more, modifications of the weed control program can be considered. Labels of materials under consideration should be checked carefully for restrictions prior to use. Some herbicides require reduced rates around young trees to minimize potential damage to resets planted among older trees. Specially modified herbicide applicators are available which enable the equipment operator to deliver reduced rates or a different herbicide mix around young trees.

To minimize herbicide contact to young trees, many growers apply a wrap or guard around the lower 12 to 16 inches of the tree trunk. When using these wraps be sure to monitor the protective structure for ants or other pests that may damage the tree trunk.

## SPROUTING

Resets require periodic sprout removal. The use of tree wraps usually reduces the need for sprout control. Wraps often stay in place for up to 3 years. They should, however, be checked periodically for the presence of ants or fungal diseases. Reduced sprouting may be enough to justify their use. There are no simple answers to the use of wraps. Each situation is different and requires careful horticultural and economic consideration to arrive at the best procedure of maintenance, inspection, and management.

## **IRRIGATION & DRAINAGE**

Young citrus trees require frequent but moderate water application for survival and proper growth. Competition for water is accentuated by nearby older trees or if weeds are allowed to grow close to the young trees. Anything that can be done to discourage competition for available water should be beneficial to the young tree. Irrigation systems should be in place before planting trees. Special modifications to the irrigation pattern by inverting the micro-sprinkler so that the surface wetting area is reduced or by increasing irrigation frequency can be good strategies to supply water for resets. However, the irrigation frequency necessary to sustain a mature grove is rarely adequate for good growth of newly-set trees, and young trees should be checked frequently to be certain they are receiving sufficient water. Drainage is as important as irrigation. Excess water must be removed from the rootzone. The concept of total water management must be practiced. If either system -irrigation or drainage- is not designed, operated, and maintained properly, then the maximum profit potential of a grove cannot be achieved. In Florida, both surface and subsoil drainage is necessary to obtain adequate root systems for the trees.

## FERTILIZATION

Reset fertilization requires an extra effort beyond the needs of the bearing grove. Frequent application of water-soluble fertilizers with irrigation water (fertigation) can increase overall fertilizer

use efficiency. If the grove is under a fertigation program, there is no need for special care in terms of nutrition for resets. Great care must be taken to ensure that proper rates of fertilizer materials are dispensed to prevent nutritional deficiencies or toxicities. Frequent light applications usually produce best results and lessen the danger of leaching but these practices need to be evaluated for cost effectiveness. The use of controlled-release fertilizers for resets is a better option than making multiple trips throughout the year to scattered resets throughout large blocks.

### PEST CONTROL

Because young trees have more frequent flushing cycles than mature trees, they are more attractive and sensitive to pests. Therefore, special care is needed to keep the citrus psyllid and leafminer under control to reduce their damage to new leaves and to reduce the severity of citrus canker and the spread of citrus greening. Relying solely on foliar contact insecticides for resets is not a good strategy. Soil-applied systemic insecticides (neonicotinoids) which provide 6-8 weeks of control are the most effective tool for managing psyllids and leafminers on resets. Currently, three neonicotinoid products are registered for use in citrus: imidacloprid (Admire, Alias, Couraze, Nuprid), thiamethoxam (Platinum), and clothianidin (Belay). Various generic formulations are also available. Resets should also benefit from foliar contact pesticides and from foliar nutrition used on mature trees.

### GROVE PLAT

Since resets are usually scattered throughout a block of much larger trees, they are often difficult to locate and may be accidentally overlooked, resulting in inadequate care. An annually updated grove plat is probably the best method for assessing general grove condition and productivity. Plats can be prepared by hand or with the assistance of a computer. This can help determine the number of trees which will be needed and where they should be placed. Reset plats can be prepared to later help equipment operators locate newly-planted trees for periodic care.

---

## **Scattered resets in a citrus grove.**



# Danger of Heat Stress

**Be alert to early warnings of heat stress, both in yourself and in your co-workers.**

**Heat stress needs to be taken seriously.**

Working in a hot environment puts stress on the body's cooling system. When heat is combined with other stresses like hard physical work, loss of fluids, or fatigue it may lead to heat-related illness. Individuals over 40 years of age need to take extra care when the weather is hot because their ability to sweat declines as they age. However, heat stress can also affect individuals who are young and fit.

## **POINTS TO EMPHASIZE:**

- Drink plenty of water to keep body fluid levels up
  - Get out of the heat occasionally
- Water is crucial to help the body adjust to high temperatures. The rate of water intake must be equal to the rate of water loss by perspiration to keep body temperature normal. **When it's hot, drink plenty of water!**

Your body must work even harder to get rid of excess heat when conditions are both hot and humid. Unfortunately, water can't evaporate as readily under muggy conditions. The process is easier if the surrounding air is moving. That's why we welcome a cool breeze, or turn on a fan when the air is "sticky".

Sickness and accident rates increase when heavy work is done at temperatures above 86 F.

Don't push yourself beyond your limits. It could be harmful to your health, and could put you at increased risk of having an accident.



## **Heat stress hazards**

1. **Heat cramps:** Heavy sweating drains the body of salt, which cannot be replaced by simply drinking water. Painful cramps occur in the arms, legs, or stomach while on the job, or later at home. Move to a cool area at once if cramping is experienced. Loosen clothing and drink cool, commercial fluid replacement beverage. Seek medical aid if the cramps are severe, or don't go away.



**2. Heat exhaustion:** Inadequate water and salt intake causes the body's cooling system to break down. Symptoms include heavy sweating, cool, moist skin, body temperature over 100 F, weak pulse, and normal or low blood pressure. The victim is likely to be tired, weak, clumsy, upset, or confused. He will be very thirsty, and will breathe rapidly. His vision may be blurred. **Get medical help immediately!** Heat exhaustion can lead to heat stroke, which can kill. Move the person to a cool, shaded area. Loosen or remove excess clothing. Provide cool, lightly-salted water. Fan and spray the victim with cool water.

**3. Heat stroke can kill a person quickly!** Once the body uses up all its water and salt, sweating ceases. Temperature can rise quickly. You can assume a person is suffering from heat stroke if their body temperature is over 105 F, and any of the following symptoms are present:

- weakness, confusion, distress, strange behavior
- hot, dry, red skin
- rapid pulse
- headache or dizziness
- In later stages of a heat stroke, a victim may pass out and have convulsions

**Call an ambulance immediately** if heat stroke is suspected. The victim's life may be on the line! Until help arrives, move the victim to a cool area and remove excess clothing. Fan and spray them with cool water. Offer sips of water if the victim is conscious.

### Heatwave guidelines

The following measures should help prevent the development of heat-related illnesses.

- Slow down in hot weather. Your body's temperature regulating system faces a much greater workload when temperature and humidity are high.

- Heed early warnings of heat stress, such as headache, heavy perspiration, high pulse rate, and shallow breathing. Take a break immediately and get to a cooler location. **Watch for heat stress signs among your co-workers.**

- Dress for hot weather. Lightweight, light-colored clothing reflects heat.

- **Drink plenty of water.** Don't let yourself "dry out".

- Try to get used to warm weather gradually. Take it easy for those first three hot days. Your body will have a better chance to adjust if you take it slow.

- Get out of the heat occasionally. Physical stress increases with time in hot weather.

Take breaks in a cool, shady location.

- Wear a hat and long-sleeved shirt to prevent burning (which can increase the risk of skin cancer.)

### "Do's" and "Don'ts" of preventing heat-related illnesses

<p><b>DO:</b>          Drink plenty of water          Take breaks in a cool, shady area          Watch for symptoms of a heat stress, both in yourself and co-workers</p>	<p><b>DON'T:</b>          Ignore symptoms of heat stress          Try to "keep up" with the rest of the crew, even though you feel ill</p>
---	--

## MANAGING HEAT STRESS

Heat stress is caused by working in hot conditions and when the body builds up more heat than it can cope with. Several factors work together to cause heat stress. Before beginning a task, think about whether any of these factors are likely to be a problem. Consider making adjustments in the task itself or in the workplace conditions, including: heat factors--temperature, humidity, air movement, and sunlight; workload--the amount of effort a task takes; drinking water intake; and scheduling.

High temperatures, high humidity, and sunlight increase the likelihood of heat stress. Air movement, from wind or from fans, may provide cooling. Because hard work causes the body to produce heat, a person is more likely to develop heat stress when working on foot than when driving a vehicle. Lifting or carrying heavy containers or equipment also increases the likelihood of overheating. Use fans, ventilation systems (indoors), and shade whenever possible. A work area or vehicle sometime can be shaded by a tarp or canopy or provided with fans or air conditioners. Consider wearing cooling clothes that help keep the body cool.

People who have become used to working in the heat are less likely to be affected by heat stress. To become adjusted to hot work environments, do about two hours of light work per day in the heat for several days in a row; then gradually increase the work period and the workload for the next several days. An adjustment period of at least seven days is recommended. If the warm weather occurs gradually, workers may adjust naturally to working in hot conditions.

Whenever it is practical, choose coveralls that allow air to pass through. Woven fabrics (cotton, or cotton-polyester blends) allow air to pass through fairly easily. Rubberized or plastic fabrics and fabrics coated with chemical-resistant barrier layers allow almost no air to pass through.

Perspiration or evaporation of sweat cools the body. Under the conditions that lead to heat stress, the body produces a large amount of sweat. Unless the water lost in sweat is replaced, body temperature will rise. Drink plenty of water before, during, and after work during heat stress conditions. Do not rely on thirst alone to guide you. A person can lose a dangerous amount of water before feeling thirsty, and the feeling of thirst may stop long before fluids are replaced. Be sure to keep body weight fairly constant. All weight lost because of sweating should be regained every day.

When the combination of temperature, sunlight, humidity, and workload is likely to lead to overheating, use scheduling to avoid heat stress. Schedule tasks requiring the heaviest workload during the coolest part of the day. When heat stress risk is high, schedule frequent breaks to allow the body to cool. Anyone who gets dangerously hot should stop work immediately and cool down. If necessary, shorten the time between breaks.

The above steps will prevent most heat stress problems. But under extremely hot conditions when cooling devices cannot be used, it may be necessary to stop work until conditions improve.

## Signs and Symptoms of Heat Stress



**Heat stress, even mild heat stress, makes people feel ill** and impairs their ability to do a good job. They may get tired quickly, feel weak, be less alert, and less able to use good judgment.

Learn the signs and symptoms of heat stress and take immediate action to cool down if you observe:

fatigue (exhaustion, muscle weakness),

headache, nausea, and chills,

dizziness and fainting,

loss of coordination,

severe thirst and dry mouth,

altered behavior (confusion, slurred speech, quarrelsome or irrational attitude).

**Severe heat stress (heat stroke) is a serious illness.** Unless victims are cooled quickly, they can die. Severe heat stress is fatal to more than 10 percent of its victims--even young, healthy adults. Victims may remain sensitive to heat for months and be unable to return to the same work.

**Heat cramps can be painful.** These are muscle spasms in the legs, arms, or stomach caused by loss of body salts through heavy sweating. To relieve cramps, drink cool water or "sports drinks." Stretching or kneading the muscles may temporarily relieve the cramps.

## First Aid for Heat Stress

It is not always easy to tell the difference between heat stress illness and pesticide poisoning. The signs and symptoms are similar. **Don't waste time trying to decide what is causing the illness. Get medical help right away.**

Get the victim into a shaded or cool area.

Cool victim as rapidly as possible by sponging or splashing skin, especially face, neck, hands, and forearms, with cool water or, when possible, immersing in cool water.

Carefully remove clothing that may be making the victim hot,


Have the victim, if conscious, drink as much cool water as possible.

Keep the victim quiet until help arrives.

**Severe heat stress (heat stroke) is a medical emergency! Cool victim immediately. Brain damage and death may result if treatment is delayed.**

# Pesticide Resistance and Resistance Management

Lauren Diepenbrock, Megan Dewdney, and Ramdas Kanissery



Populations of animals, fungi, bacteria, and plants possess the ability to respond to sustained changes or stresses in their environment in ways that enable the continued survival of the species. Such environmental stresses include physical factors (e.g., temperature or humidity), biological factors (e.g., predators, parasites, or pathogens) and environmental contaminants. In any population, a small percentage of individuals will be better able to respond to new stresses because of unique traits or characteristics that they possess. Consequently, those individuals will survive, reproduce, and become more common in a population. This phenomenon is commonly referred to as "survival of the fittest."

Many pest species, such as the citrus rust mite, are exceptionally well-equipped to respond to environmental stresses because of their short generation time and large reproductive potential. The use of chemical sprays to control insect, mite, and fungal diseases of citrus creates a potent environmental stress. There are now many examples of pests and pathogens that have responded by developing resistance to one or more pesticides. Pesticide-resistant individuals are those that have developed the ability to tolerate doses of a toxicant that would be lethal to the majority of individuals. The resistance mechanisms can vary according to pest species and/or the class of chemical to which the pest is exposed. Resistance mechanisms include an increased capacity to detoxify the pesticide once it has entered the pest's body, a decreased sensitivity of the target site that the pesticide acts upon, a decreased penetration of the pesticide through the cuticle, or sequestration of the pesticide within the organism. The main resistance mechanism for pathogens is a change in the target site so that the pathogen is less susceptible or fully resistant. A single resistance mechanism can sometimes provide defense against different classes of chemicals and this is known as *cross-resistance*. When more than one resistance mechanism is expressed in the same individual, this individual is said to show *multiple resistance*.

Because the traits for resistance are passed from one generation to the next, continued stress from a pesticide may, over time, create resistance in the majority of individuals in a population. From an operational perspective, this process would be expressed as a gradual decrease and eventual loss of effectiveness of a chemical. Resistance to a particular chemical may be stable or unstable. When resistance is stable, the pest population does not revert to a susceptible state even if the use of that chemical is discontinued. When resistance is unstable and use of the chemical is temporarily discontinued, the population will eventually return to a susceptible state, at which time the

chemical in question could again be used to manage that pest. However, in this situation, previously resistant populations may eventually show resistance again.

Of the factors that affect the development of resistance—which include the pest's or pathogen's biology, ecology and genetics—only the operational factors can be manipulated by the grower. The key operational factor that will delay the onset of pesticidal resistance and prolong the effective life of a compound is to assure the survival of some susceptible individuals to dilute the population of resistant individuals. The following operational procedures should be on a grower's checklist to steward sound pesticidal resistance management for acaricides, insecticides, fungicides, and herbicides:

Never rely on a single pesticide class.

Integrate chemical control with effective, complementary cultural and biological control practices.

Always use pesticides at recommended rates and strive for thorough coverage.

When there is more than one generation of pest, alternate different pesticide classes.

Do not use tank mixtures of products that have the same mode of action. If control with a pesticide fails, do not re-treat with a chemical that has the same mode of action.

Reports of resistance have been documented for certain acaricides used to control citrus rust mite and fungicides used to combat diseases in Florida. Resistance to Benlate developed in the greasy spot fungus shortly after the product was introduced about 30 years ago and is still widespread. Benlate resistance also occurs in the scab fungus in isolated situations and is stable. Resistance has been detected in tangerine groves with Alternaria brown spot to strobilurin fungicides (Abound, Gem, and Headline and contained in the mixtures Pristine and Quadris Top) but no resistance has developed to ferbam. Dicofol resistance in citrus rust mite was detected throughout the citrus industry about 10 years ago, but resistance proved to be unstable and usage of dicofol has continued. Agri-mek tolerance in citrus rust mite is of concern and growers should follow sound resistant management practices when using this product.

The following tables are provided to aid in the rotation of pesticides with different modes of action within a season or from year to year. There is a separate table for insecticides/acaricides, fungicides, and herbicides. The information in these tables was derived from information produced by the Insecticide Resistance Action Committee (IRAC) (<http://www.iraconline.org/>), Fungicide Resistance Action Committee (FRAC) (<http://www.frac.info/>), and the Herbicide Resistance Action Committee (HRAC) (<http://hracglobal.com/pages/classificationofherbicidesiteofaction.aspx>). Each table lists the number (or letter in the case of herbicides) of the group code for each pesticide class, the group name or general description of that group of pesticides, the common name of pesticides used in citrus production that belong to each group, and examples of trade names of pesticides for each common name listed. When using the table to rotate between using products with different modes of action, choose products with a different group code than previously used in the grove during the current growing season. In the case of insecticides/acaricides, many of these pesticides are broken into subgroups. It is unclear whether cross resistance will occur between these subgroups. When possible, it is recommended to rotate with an entirely different group. (Note: The IRAC and FRAC mode of action systems both use a similar numbering system. There is no cross-resistance potential between the insecticides and fungicides.) Products with broad-based activity such as sulfur, copper, and oil are not included in this list because the development of resistance to them is not likely.

For more details, go to: <https://edis.ifas.ufl.edu/pdf/CG/CG026/CG026-Dfryt7p45d.pdf>

# Citrus Spray Programs

Dr. Jawwad Qureshi and Dr. Phil Stansly, UF IFAS- Immokalee

Asian citrus psyllid (ACP) control has been the main objective of Florida citrus growers for more than 10 years. While some may question the value of controlling ACP in trees with high HLB incidence, replicated field studies have shown the economic benefit of maintaining young flush pathogen free. Good ACP control starts with effective dormant sprays that will control ACP when populations are low, reducing ACP infestation and thus HLB infection of the all-important spring flush. Pyrethroids (Danitol, Baythroid or Mustang) and organophosphates (dimethoate, chlorpyrifos, or Imidan) provide great winter season control of ACP. Best not to use pyrethroids or OPs again during the year except for border sprays which will reduce the need for whole block applications. Follow up with bloom sprays of labeled products to clean up stragglers. Subsequent whole block sprays should target ACP as well as other pests like rust mites and leafminers that may be problematic.

The table below offers alternative products for different months, depending on which pests are of major concern at the time. Neonicotinoids like imidacloprid, thiamethoxam or clothianidin have not been included as spray options due to their importance for controlling ACP in young trees. Superscripts after the pesticide name are now in sequential order to facilitate use and correspond to superscripts after pests controlled. Make choices based on: (1) effectiveness against ACP and other pests that may be problematic, (2) avoiding repetition of any insecticide mode of action in the interest of resistance management, and (3) rebuilding and maintaining an effective natural enemy complex in the grove. Confining the broad-spectrum insecticides (pyrethroids and organo-phosphates) to the winter season and border sprays during growing season will help conserve these products as well as populations of beneficial insects and mites.

## Spray Options for Citrus Pest Management

**Dormant Season**

**Growing Season**

Months	Nov-Dec	Jan	Feb-Mar	Apr	May - June	July - Aug	Sep-Oct
<b>Products</b>  <b>* Labeled for bloom</b>	OP <sup>1</sup> (e.g. Imidan, Dimethoate, chlorpyrifos)	Pyrethroid <sup>2</sup> (Mustang, Danitol, Baythroid)	*Sivanto <sup>3</sup> *Movento <sup>4</sup> *Portal <sup>5</sup> *Micromite <sup>6</sup> Intrepid <sup>7</sup> Exirel <sup>8</sup>	Portal <sup>5</sup> Micromite <sup>6</sup> Exirel <sup>8</sup> Apta <sup>9</sup> Sivanto <sup>3</sup> Oil <sup>13</sup>	Movento <sup>4</sup> Delegate <sup>11</sup> Abamectin <sup>12</sup> Knack <sup>14</sup> Exirel <sup>8</sup> Apta <sup>9</sup> Sivanto <sup>3</sup> Oil <sup>13</sup> MinectoPro <sup>10</sup>	Sivanto <sup>3</sup> Apta <sup>9</sup> OP <sup>1</sup> MinectoPro <sup>10</sup> Oil <sup>13</sup>	Movento <sup>4</sup> Delegate <sup>11</sup> Apta <sup>9</sup> Sivanto <sup>3</sup> Oil <sup>13</sup>
<b>Pests</b>	ACP Weevils	ACP Weevils	ACP Mites Leafminer Weevils Scales Aphids	ACP Mites Leafminer Weevils Aphids	ACP Rustmite Leafminer Scales	ACP	ACP Rustmite Leafminer
<p><b>ACP+++<sup>1,2,3,4,8,9,10</sup> ACP++<sup>5,11</sup> ACP+<sup>6,12</sup> Leafminer<sup>6,7,8,9,11,12</sup> Rustmite<sup>4,12</sup> Scales<sup>4,13</sup> Aphids<sup>3,4</sup> Mealybugs<sup>3,4</sup> (+++ excellent, ++ good, + fair)</b></p>							

**Flatwoods Citrus newsletter** by regular mail stopped this month.

You will receive your copy only through e-mail or through the following link:

<https://citrusagents.ifas.ufl.edu/newsletters/>

If you did not receive the *Flatwoods Citrus* newsletter and would like to be on our e-mailing list, please check this box and complete the information requested below.

If you wish to be removed from our mailing list, please check this box and complete the information requested below.

Please send: Dr. Mongi Zekri  
Multi-County Citrus Agent  
Hendry County Extension Office  
P.O. Box 68  
LaBelle, FL 33975  
or E-mail: [maz@ufl.edu](mailto:maz@ufl.edu)

---

Subscriber's Name: \_\_\_\_\_

Company: \_\_\_\_\_

Phone: \_\_\_\_\_

E-mail: \_\_\_\_\_

---

**Racial-Ethnic Background**

American Indian or native Alaskan

White, non-Hispanic

Asian American

Black, non-Hispanic

Hispanic

**Gender**

Female

Male